



09/284,534
Part of Paper #7

Kongeriget Danmark

Patent application No.: PA 2000 00291
Date of filing: 24 February 2000
Applicant: Novozymes A/S
Krogshøjvej 36
DK-2880 Bagsværd

This is to certify the correctness of the following information:

The attached photocopy is a true copy of the following document:

- The specification, sequence listing, claims and abstract as filed with the application on the filing date indicated above.

By assignment dated 17 Nov 2000 and filed on 01 Dec 2000, the application has been assigned to Novozymes A/S



Patent- og
Varemærkestyrelsen
Erhvervsministeriet

TAASTRUP 22 May 2001

Lizzi Vester

Lizzi Vester
Head of Section

24 FEB. 2000

FAMILY 44 XYLOGLUCANASES

Modtaget

The present invention relates to xyloglucanases belonging to family 44 of glycosyl hydrolases, preferably to enzymes exhibiting xyloglucanase activity as their major enzymatic activity in the neutral and alkaline pH ranges; to a method of producing such enzymes; and to methods for using such enzymes in the textile, detergent and cellulose fiber processing industries.

10

BACKGROUND OF THE INVENTION

Xyloglucan is a major structural polysaccharide in the primary (growing) cell wall of plants. Structurally, xyloglucans consists of a cellulose-like beta-1,4-linked glucose backbone which is frequently substituted with various side chains. The xyloglucans of most dicotyledonous plants, some monocotyledons and gymnosperms are highly branched polysaccharides in which approx. 75% of the glucose residues in the backbone bear a glycosyl side chain at O-6. The glycosyl residue that is directly attached to the branched glucose residue is invariably alfa-D-xylose. Up to 50% of the side chains in the xyloglucans contain more than one residue due to the presence of beta-D-galactose or alfa-L-fucose-(1-2)-beta-D-galactose moieties at O-2 of the xylose residues (C. Ohsumi and T. Hayashi (1994) Plant and Cell Physiology 35:963-967; G. J. McDougall and S. C. Fry (1994) Journal of Plant Physiology 143:591-595; J. L. Acebes et al. (1993) Phytochemistry 33:1343-1345). On acid hydrolysis, the xyloglucan extracted from cotton fibers yielded glucose, xylose, galactose and fucose in the ratio of 50:29:12:7 (Hayashi et al., 1988).

Xyloglucans produced by solanaceous plants are unusual in that typically only 40% of the beta-1,4-linked glucose residues bear a glycosyl side chain at O-6. Furthermore, up to 60% of the xylose residues are substituted at O-2 with alfa-L-arabinose residues and some solanaceous plants, such as potato, also have xyloglucans with beta-D-galactose substituents at O-2 of some of the xylose residues (York et al (1996)).

Xyloglucan is believed to function in the primary wall of plants by crosslinking cellulose-microfibrils, forming a

cellulose-xyloglucan network. This network is considered necessary for the structural integrity of primary cell-walls (Carpita et al., 1993). Another important function of xyloglucan is to act as a repository for xyloglucan subunit oligo-
5 saccharides that are physiologically active regulators of plant cell growth. Xyloglucan subunits may also modulate the action of a xyloglucan endotransglycosylase (XET), a cell-wall associated enzyme that has been hypothesized to play a role in the elongation of plant cell walls. Therefore xyloglucan might play
10 an important role in wall loosening and consequently cell expansion (Fry et al., 1992).

The seeds of many dicotyledonous species contain xyloglucan as the major polysaccharide storage reserve. This type of xyloglucan, which is localized in massive thickenings on the
15 inside of the seed cotyledon cell wall, is composed mainly of glucose, xylose and galactose (Rose et al., 1996).

Seeds of the tamarind tree *Tamarindus indica* became a commercial source of gum in 1943 when the gum was found useful as a paper and textile size. Sizing of jute and cotton with
20 tamarind xyloglucan has been extensively practiced in Asia owing to the low cost of the gum and to its excellent properties. Food applications of tamarind xyloglucan include use in confections, jams and jellies and as a stabilizer in ice cream and mayonnaise (Whistler et al., 1993).

25 Xyloglucanase activity is not included in the classification of enzymes provided by the Enzyme Nomenclature (1992). Hitherto, this enzymatic activity has simply been classified as glucanase activity and has often been believed to be identical to cellulolytic activity (EC 3.2.1.4), i.e.
30 activity against β -1,4-glycosidic linkages in cellulose or cellulose derivative substrates, or at least to be a side activity in enzymes having cellulolytic activity. However, a true xyloglucanase is a true xyloglucan specific enzyme capable of catalyzing the solubilisation of xyloglucan to xyloglucan
35 oligosaccharides but which does not exhibit substantial cellulolytic activity, e.g. activity against the conventionally used cellulose-like substrates CMC (carboxymethylcellulose), HE cellulose and Avicel (microcrystalline cellulose). A xyloglucanase cleaves the beta-1,4-glycosidic linkages in the

backbone of xyloglucan.

Xyloglucanase activity is described by Vincken et al. (1997) who characterizes three different endoglucanases from *Trichoderma viride* (similar to *T. reesei*) which all have high activity against cellulose or CMC and show that the EndoI (belonging to family 5 of glycosyl hydrolases, see Henrissat, B. et al. (1991, 1993)) has essentially no (i.e. very little) activity against xyloglucan, and that EndoV (belonging to the family 7 of glycosyl hydrolases) and EndoIV (belonging to the family 12 of glycosyl hydrolases) both have activity against xyloglucan and CMC, respectively, of the same order of magnitude.

International Patent Publication WO 94/14953 discloses a family 12 xyloglucanase (EG II) cloned from the fungus *Aspergillus aculeatus* and expressed in the fungus *Aspergillus oryzae*.

International Patent Publication WO 99/02663 discloses xyloglucanases cloned from *Bacillus licheniformis* (family 12) and *Bacillus agaradhaerens* (family 5) and expressed in *Bacillus subtilis*.

It is an object of the present invention to provide an enzyme with a high xyloglucanase activity at an alkaline pH which xyloglucanase exhibits excellent performance in conventional detergent compositions.

SUMMARY OF THE INVENTION

The inventors have now found enzymes having substantial xyloglucanase activity, which enzymes belong to family 44 of glycosylhydrolases and exhibit excellent performance in conventional detergent compositions, especially liquid detergent compositions.

Accordingly, the present invention relates to an enzyme preparation comprising a xyloglucanase belonging to family 44 of glycosylhydrolases and exhibiting a relative activity of at least 30% between pH 5.0 and pH 8.0.

The inventors have also succeeded in cloning and expressing a family 44 xyloglucanase, i.e. the invention relates in further aspects to a family 44 xyloglucanase which is (a) a polypeptide encoded by the DNA sequence of positions 121-1677 of SEQ ID

NO:1, (b) a polypeptide produced by culturing a cell comprising the sequence of SEQ ID NO:1 under conditions wherein the DNA sequence is expressed, (c) a xyloglucanase enzyme having a sequence of at least 60% identity to positions 40-559 of SEQ ID NO:2 when identity is determined by GAP provided in the GCG program package using a GAP creation penalty of 3.0 and GAP extension penalty of 0.1, or (d) a polypeptide encoded by the xyloglucanase encoding part of the DNA sequence obtainable from the plasmid in *Escherichia coli* DSM 13321; and to a family 44 xyloglucanase which is (a) a polypeptide encoded by the DNA sequence of positions 121-1677 of SEQ ID NO:3, (b) a polypeptide produced by culturing a cell comprising the sequence of SEQ ID NO:3 under conditions wherein the DNA sequence is expressed, or (c) a polypeptide encoded by the xyloglucanase encoding part of the DNA sequence obtainable from the plasmid in *Escherichia coli* DSM 13322; and to a family 44 xyloglucanase which is (a) a polypeptide encoded by the DNA sequence of positions 121-1677 of SEQ ID NO:5, (b) a polypeptide produced by culturing a cell comprising the sequence of SEQ ID NO:5 under conditions wherein the DNA sequence is expressed, or (c) a polypeptide encoded by the xyloglucanase encoding part of the DNA sequence obtainable from the plasmid in *Escherichia coli* DSM 13323; and to an isolated polynucleotide molecule encoding a polypeptide having xyloglucanase activity which polynucleotide molecule hybridizes to a denatured double-stranded DNA probe under medium stringency conditions, wherein the probe is selected from the group consisting of DNA probes comprising the sequence shown in positions 121-1677 of SEQ ID NO:1, 3 or 5, and DNA probes comprising a subsequence of positions 121-1677 of SEQ ID NO:1, 3 or 5, the subsequence having a length of at least about 100 base pairs.

In further aspects, the invention provides an expression vector comprising a DNA segment which is e.g. a polynucleotide molecule of the invention; a cell comprising the DNA segment or the expression vector; and a method of producing a exhibiting xyloglucanase enzyme, which method comprises culturing the cell under conditions permitting the production of the enzyme, and recovering the enzyme from the culture.

In yet another aspect the invention provides an isolated

xyloglucanase enzyme characterized in (i) being free from homologous impurities and (ii) being produced by the method described above.

The novel enzyme of the present invention is useful for the treatment of cellulosic material, especially cellulose-containing fiber, yarn, woven or non-woven fabric. The treatment can be carried out during the processing of cellulosic material into a material ready for garment manufacture or fabric manufacture, e.g. in the desizing or scouring step; or during industrial or household laundering of such fabric or garment.

Accordingly, in further aspects the present invention relates to a detergent composition comprising a xyloglucanase enzyme having substantial xyloglucanase activity in the neutral or alkaline range; and to use of the enzyme of the invention for the treatment of cellulose-containing fibers, yarn, woven or non-woven fabric.

The present invention has now made it possible to use a xyloglucanase in detergent compositions for removing or bleaching certain soils or stains present on laundry, especially soils and spots resulting from xyloglucan-containing food, plants, and the like. Further, it is contemplated that treatment with detergent compositions comprising the novel enzyme can prevent binding of certain soils to the xyloglucan left on the cellulosic material.

25

DETAILED DESCRIPTION OF THE INVENTION

MICROBIAL SOURCES

For the purpose of the present invention the term "obtained from" or "obtainable from" as used herein in connection with a specific source, means that the enzyme is produced or can be produced by the specific source, or by a cell in which a gene from the source have been inserted.

It is at present contemplated that the xyloglucanase of the invention may be obtained from a gram positive bacterium belonging to a strain of the genus *Bacillus*, in particular a strain of *Paenibacillus*.

In a preferred embodiment, the xyloglucanase of the invention is obtained from the strain *Paenibacillus polymyxa*,

ATCC 832, which is publicly available from American Type Culture Collection (ATCC). This is the type strain of *Paenibacillus polymyxa*. It is at present contemplated that a DNA sequence encoding an enzyme with an amino acid sequence identity of at least 60% to the enzyme of the invention may be obtained from other strains belonging to the genus *Paenibacillus*.

Further, the strain *Paenibacillus sp.* was deposited by the inventors according to the Budapest Treaty on the International Recognition of the Deposit of Microorganisms for the Purposes of Patent Procedure at the Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH, Mascheroder Weg 1b, D-38124 Braunschweig, Federal Republic of Germany, on 18 February 2000 under the deposition number DSM 13329.

A plasmid comprising a DNA sequence encoding a xyloglucanase of the invention has been transformed into a strain of the *Escherichia coli* which was deposited by the inventors according to the Budapest Treaty on the International Recognition of the Deposit of Microorganisms for the Purposes of Patent Procedure at the Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH, Mascheroder Weg 1b, D-38124 Braunschweig, Federal Republic of Germany, on 16 February 2000 under the deposition number DSM 13321. It is contemplated that the DNA sequence of this plasmid comprises the DNA sequence of SEQ ID NO:1.

A plasmid comprising a DNA sequence encoding a xyloglucanase of the invention has been transformed into a strain of the *Escherichia coli* which was deposited by the inventors according to the Budapest Treaty on the International Recognition of the Deposit of Microorganisms for the Purposes of Patent Procedure at the Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH, Mascheroder Weg 1b, D-38124 Braunschweig, Federal Republic of Germany, on 16 February 2000 under the deposition number DSM 13322. It is contemplated that the DNA sequence of this plasmid comprises the DNA sequence of SEQ ID NO:3.

A plasmid comprising a DNA sequence encoding a xyloglucanase of the invention has been transformed into a strain of the *Escherichia coli* which was deposited by the inventors according to the Budapest Treaty on the International

Recognition of the Deposit of Microorganisms for the Purposes of Patent Procedure at the Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH, Mascheroder Weg 1b, D-38124 Braunschweig, Federal Republic of Germany, on 16 February 2000 under the
5 deposition number DSM 13323. It is contemplated that the DNA sequence of this plasmid comprises the DNA sequence of SEQ ID NO:5.

DEFINITIONS

10 In the present context, the term "enzyme preparation" is intended to mean either be a conventional enzymatic fermentation product, possibly isolated and purified, from a single species of a microorganism, such preparation usually comprising a number of different enzymatic activities; or a mixture of monocomponent
15 enzymes, preferably enzymes derived from bacterial or fungal species by using conventional recombinant techniques, which enzymes have been fermented and possibly isolated and purified separately and which may originate from different species, preferably fungal or bacterial species; or the fermentation
20 product of a microorganism which acts as a host cell for expression of a recombinant xyloglucanase, but which microorganism simultaneously produces other enzymes, e.g. xyloglucanases, proteases, or cellulases, being naturally occurring fermentation products of the microorganism, i.e. the
25 enzyme complex conventionally produced by the corresponding naturally occurring microorganism.

In the present context the term "expression vector" denotes a DNA molecule, linear or circular, that comprises a segment encoding a polypeptide of interest operably linked to additional
30 segments that provide for its transcription. Such additional segments may include promoter and terminator sequences, and may optionally include one or more origins of replication, one or more selectable markers, an enhancer, a polyadenylation signal, and the like. Expression vectors are generally derived from
35 plasmid or viral DNA, or may contain elements of both. The expression vector of the invention may be any expression vector that is conveniently subjected to recombinant DNA procedures, and the choice of vector will often depend on the host cell into which the vector is to be introduced. Thus, the vector may be an

autonomously replicating vector, i.e. a vector which exists as an extrachromosomal entity, the replication of which is independent of chromosomal replication, e.g. a plasmid. Alternatively, the vector may be one which, when introduced into a host cell, is integrated into the host cell genome and replicated together with the chromosome(s) into which it has been integrated.

The term "recombinant expressed" or "recombinantly expressed" used herein in connection with expression of a polypeptide or protein is defined according to the standard definition in the art. Recombinant expression of a protein is generally performed by using an expression vector as described immediately above.

The term "isolated", when applied to a polynucleotide molecule, denotes that the polynucleotide has been removed from its natural genetic milieu and is thus free of other extraneous or unwanted coding sequences, and is in a form suitable for use within genetically engineered protein production systems. Such isolated molecules are those that are separated from their natural environment and include cDNA and genomic clones. Isolated DNA molecules of the present invention are free of other genes with which they are ordinarily associated, but may include naturally occurring 5' and 3' untranslated regions such as promoters and terminators. The identification of associated regions will be evident to one of ordinary skill in the art (see for example, Dynan and Tijan, Nature 316:774-78, 1985). The term "an isolated polynucleotide" may alternatively be termed "a cloned polynucleotide".

When applied to a protein/polypeptide, the term "isolated" indicates that the protein is found in a condition other than its native environment. In a preferred form, the isolated protein is substantially free of other proteins, particularly other homologous proteins (i.e. "homologous impurities" (see below)). It is preferred to provide the protein in a greater than 40% pure form, more preferably greater than 60% pure form.

Even more preferably it is preferred to provide the protein in a highly purified form, i.e., greater than 80% pure, more preferably greater than 95% pure, and even more preferably greater than 99% pure, as determined by SDS-PAGE.

The term "isolated protein/polypeptide" may alternatively be termed "purified protein/polypeptide".

The term "homologous impurities" means any impurity (e.g. another polypeptide than the polypeptide of the invention) which originate from the homologous cell where the polypeptide of the invention is originally obtained from.

The term "obtained from" as used herein in connection with a specific microbial source, means that the polynucleotide and/or polypeptide produced by the specific source, or by a cell in which a gene from the source have been inserted.

The term "operably linked", when referring to DNA segments, denotes that the segments are arranged so that they function in concert for their intended purposes, e.g. transcription initiates in the promoter and proceeds through the coding segment to the terminator

The term "polynucleotide" denotes a single- or double-stranded polymer of deoxyribonucleotide or ribonucleotide bases read from the 5' to the 3' end. Polynucleotides include RNA and DNA, and may be isolated from natural sources, synthesized *in vitro*, or prepared from a combination of natural and synthetic molecules.

The term "complements of polynucleotide molecules" denotes polynucleotide molecules having a complementary base sequence and reverse orientation as compared to a reference sequence. For example, the sequence 5' ATGCACGGG 3' is complementary to 5' CCCGTGCAT 3'.

The term "degenerate nucleotide sequence" denotes a sequence of nucleotides that includes one or more degenerate codons (as compared to a reference polynucleotide molecule that encodes a polypeptide). Degenerate codons contain different triplets of nucleotides, but encode the same amino acid residue (i.e., GAU and GAC triplets each encode Asp).

The term "promoter" denotes a portion of a gene containing DNA sequences that provide for the binding of RNA polymerase and initiation of transcription. Promoter sequences are commonly, but not always, found in the 5' non-coding regions of genes.

The term "secretory signal sequence" denotes a DNA sequence that encodes a polypeptide (a "secretory peptide") that, as a component of a larger polypeptide, directs the larger

polypeptide through a secretory pathway of a cell in which it is synthesized. The larger peptide is commonly cleaved to remove the secretory peptide during transit through the secretory pathway.

5

POLYNUCLEOTIDES

Within preferred embodiments of the invention an isolated polynucleotide of the invention will hybridize to similar sized regions of SEQ ID NO:1, SEQ ID NO:3 or SEQ ID NO:5, or a
10 sequence complementary thereto, under at least medium stringency conditions.

In particular polynucleotides of the invention will hybridize to a denatured double-stranded DNA probe comprising either the full sequence shown in SEQ ID NO:1 or the sequence
15 shown in positions 121-1677 of SEQ ID NO:1 or the full sequence shown in SEQ ID NO:3 or the sequence shown in positions 121-1677 of SEQ ID NO:3 or the partial sequence shown in SEQ ID NO:5 or the sequence shown in positions 121-1677 of SEQ ID NO:5 or any probe comprising a subsequence of SEQ ID NO:5 or SEQ ID NO:3 or
20 SEQ ID NO:1 having a length of at least about 100 base pairs under at least medium stringency conditions, but preferably at high stringency conditions as described in detail below.

Suitable experimental conditions for determining hybridization at medium or high stringency between a nucleotide probe and a
25 homologous DNA or RNA sequence involve presoaking of the filter containing the DNA fragments or RNA to hybridize in 5 x SSC (Sodium chloride/Sodium citrate, Sambrook et al. 1989) for 10 min, and prehybridization of the filter in a solution of 5 x SSC, 5 x Denhardt's solution (Sambrook et al. 1989), 0.5 % SDS
30 and 100 µg/ml of denatured sonicated salmon sperm DNA (Sambrook et al. 1989), followed by hybridization in the same solution containing a concentration of 10ng/ml of a random-primed (Feinberg, A. P. and Vogelstein, B. (1983) Anal. Biochem. 132:6-13), 32P-dCTP-labeled (specific activity higher than 1 x 10⁹
35 cpm/µg) probe for 12 hours at ca. 45°C. The filter is then washed twice for 30 minutes in 2 x SSC, 0.5 % SDS at least 60°C (medium stringency), still more preferably at least 65°C (medium/high stringency), even more preferably at least 70°C (high stringency), and even more preferably at least 75°C (very

high stringency).

Molecules to which the oligonucleotide probe hybridizes under these conditions are detected using a x-ray film.

As previously noted, the isolated polynucleotides of the present invention include DNA and RNA. Methods for isolating DNA and RNA are well known in the art. DNA and RNA encoding genes of interest can be cloned in Gene Banks or DNA libraries by means of methods known in the art.

Polynucleotides encoding polypeptides having endogucanase activity of the invention are then identified and isolated by, for example, hybridization or PCR.

The present invention further provides counterpart polypeptides and polynucleotides from different bacterial strains (orthologs or paralogs). Of particular interest are xyloglucanase polypeptides from gram-positive alkalophilic strains, including species of *Bacillus*. Of special interest are xyloglucanase peptides from strains which are very closely related to the strain *Paenibacillus polymyxa*, ATCC 832, which is the type strain of *Paenibacillus polymyxa*.

Species homologues of a polypeptide with xyloglucanase activity of the invention can be cloned using information and compositions provided by the present invention in combination with conventional cloning techniques. For example, a DNA sequence of the present invention can be cloned using chromosomal DNA obtained from a cell type that expresses the protein. Suitable sources of DNA can be identified by probing Northern blots with probes designed from the sequences disclosed herein. A library is then prepared from chromosomal DNA of a positive cell line. A DNA sequence of the invention encoding an polypeptide having xyloglucanase activity can then be isolated by a variety of methods, such as by probing with probes designed from the sequences disclosed in the present specification and claims or with one or more sets of degenerate probes based on the disclosed sequences. A DNA sequence of the invention can also be cloned using the polymerase chain reaction, or PCR (Mullis, U.S. Patent 4,683,202), using primers designed from the sequences disclosed herein. Within an additional method, the DNA library can be used to transform or transfect host cells, and expression of the DNA of interest can be detected with an

antibody (monoclonal or polyclonal) raised against the xyloglucanase cloned from *Paenibacillus polymyxa*, e.g. from the type strain deposited as ATCC 832, or from *Paenibacillus sp.*, DSM 13329, expressed and purified as described in Materials and Methods and Examples 1, 2 and 3, or by an activity test relating to a polypeptide having xyloglucanase activity.

POLYPEPTIDES:

The sequence of amino acids nos. 40-559 of SEQ ID NO: 2, 4 and 6, respectively, is a mature xyloglucanase sequence of the catalytic active domain. The mature xyloglucanase sequence may in theory start at position 35 or 36 but the enzyme as such has the amino acid sequence starting at position 40 due to proteolytic maturing. Further, it should be noted that the genes disclosed herein (SEQ ID NOS: 1, 3, 5) *in toto* encodes multidomain enzymes, i.e. encodes a xyloglucanase domain (catalytic active domain), a mannanase domain (catalytic active domain) and a binding domain. However, only the part of the genes encoding the xyloglucanase domain is relevant for the present invention.

The present invention also provides xyloglucanase polypeptides that are substantially homologous to the polypeptide of amino acids nos. 40-559 of SEQ ID NO:2, SEQ ID NO:4 or SEQ ID NO:6 and species homologs (paralogs or orthologs) thereof. The term "substantially homologous" is used herein to denote polypeptides having 60%, preferably at least 70%, more preferably at least 75%, more preferably at least 80%, more preferably at least 85%, and even more preferably at least 90%, sequence identity to the sequence shown in amino acids nos. 40-559 of SEQ ID NO:2, 4 or 6 or their orthologs or paralogs. Such polypeptides will more preferably be at least 95% identical, and most preferably 98% or more identical to the sequence shown in amino acids nos. 40-559 of SEQ ID NO:2, 4 or 6 or its orthologs or paralogs. Percent sequence identity is determined by conventional methods, by means of computer programs known in the art such as GAP provided in the GCG program package (Program Manual for the Wisconsin Package, Version 8, August 1994, Genetics Computer Group, 575 Science Drive, Madison, Wisconsin, USA 53711) as disclosed in Needleman, S.B. and Wunsch, C.D.,

(1970), Journal of Molecular Biology, 48, 443-453, which is hereby incorporated by reference in its entirety. GAP is used with the following settings for polypeptide sequence comparison: GAP creation penalty of 3.0 and GAP extension penalty of 0.1.

- 5 The following sequence identity was found for the appended SEQ ID NOS: 2, 4 and 6:

	SEQ ID #2	SEQ ID #4	SEQ ID #6
SEQ ID #2	100%	92%	84%

10

Sequence identity of polynucleotide molecules is determined by similar methods using GAP with the following settings for DNA sequence comparison: GAP creation penalty of 5.0 and GAP extension penalty of 0.3.

- 15 Substantially homologous proteins and polypeptides are characterized as having one or more amino acid substitutions, deletions or additions. These changes are preferably of a minor nature, that is conservative amino acid substitutions (see Table 2) and other substitutions that do not significantly affect the
- 20 folding or activity of the protein or polypeptide; small deletions, typically of one to about 30 amino acids; and small amino- or carboxyl-terminal extensions, such as an amino-terminal methionine residue, a small linker peptide of up to about 20-25 residues, or a small extension that facilitates
- 25 purification (an affinity tag), such as a poly-histidine tract, protein A (Nilsson et al., EMBO J. 4:1075, 1985; Nilsson et al., Methods Enzymol. 198:3, 1991. See, in general Ford et al., Protein Expression and Purification 2: 95-107, 1991, which is incorporated herein by reference. DNAs encoding affinity tags
- 30 are available from commercial suppliers (e.g., Pharmacia Biotech, Piscataway, NJ; New England Biolabs, Beverly, MA).

However, even though the changes described above preferably are of a minor nature, such changes may also be of a larger nature such as fusion of larger polypeptides of up to 300 amino

35 acids or more both as amino- or carboxyl-terminal extensions to a polypeptide of the invention having xyloglucanase activity.

Table 1

Conservative amino acid substitutions

	Basic:	arginine
		lysine
		histidine
5	Acidic:	glutamic acid
		aspartic acid
	Polar:	glutamine
		asparagine
	Hydrophobic:	leucine
10		isoleucine
		valine
	Aromatic:	phenylalanine
		tryptophan
		tyrosine
	Small:	glycine
15		alanine
		serine
		threonine
		methionine

20 In addition to the 20 standard amino acids, non-standard amino acids (such as 4-hydroxyproline, 6-N-methyl lysine, 2-aminoisobutyric acid, isovaline and α -methyl serine) may be substituted for amino acid residues of a polypeptide according to the invention. A limited number of non-conservative amino
25 acids, amino acids that are not encoded by the genetic code, and unnatural amino acids may be substituted for amino acid residues. "Unnatural amino acids" have been modified after protein synthesis, and/or have a chemical structure in their side chain(s) different from that of the standard amino acids.
30 Unnatural amino acids can be chemically synthesized, or preferably, are commercially available, and include pipecolic acid, thiazolidine carboxylic acid, dehydroproline, 3- and 4-methylproline, and 3,3-dimethylproline.

Essential amino acids in the xyloglucanase polypeptides of
35 the present invention can be identified according to procedures known in the art, such as site-directed mutagenesis or alanine-scanning mutagenesis (Cunningham and Wells, Science 244: 1081-1085, 1989). In the latter technique, single alanine mutations are introduced at every residue in the molecule, and the

resultant mutant molecules are tested for biological activity (i.e xyloglucanase activity) to identify amino acid residues that are critical to the activity of the molecule. See also, Hilton et al., J. Biol. Chem. 271:4699-4708, 1996. The active site of the enzyme or other biological interaction can also be determined by physical analysis of structure, as determined by such techniques as nuclear magnetic resonance, crystallography, electron diffraction or photoaffinity labeling, in conjunction with mutation of putative contact site amino acids. See, for example, de Vos et al., Science 255:306-312, 1992; Smith et al., J. Mol. Biol. 224:899-904, 1992; Wlodaver et al., FEBS Lett. 309:59-64, 1992. The identities of essential amino acids can also be inferred from analysis of homologies with polypeptides which are related to a polypeptide according to the invention.

Multiple amino acid substitutions can be made and tested using known methods of mutagenesis, recombination and/or shuffling followed by a relevant screening procedure, such as those disclosed by Reidhaar-Olson and Sauer (Science 241:53-57, 1988), Bowie and Sauer (Proc. Natl. Acad. Sci. USA 86:2152-2156, 1989), WO95/17413, or WO 95/22625. Briefly, these authors disclose methods for simultaneously randomizing two or more positions in a polypeptide, or recombination/shuffling of different mutations (WO95/17413, WO95/22625), followed by selecting for functional a polypeptide, and then sequencing the mutagenized polypeptides to determine the spectrum of allowable substitutions at each position. Other methods that can be used include phage display (e.g., Lowman et al., Biochem. 30:10832-10837, 1991; Ladner et al., U.S. Patent No. 5,223,409; Huse, WIPO Publication WO 92/06204) and region-directed mutagenesis (Derbyshire et al., Gene 46:145, 1986; Ner et al., DNA 7:127, 1988).

Mutagenesis/shuffling methods as disclosed above can be combined with high-throughput, automated screening methods to detect activity of cloned, mutagenized polypeptides in host cells. Mutagenized DNA molecules that encode active polypeptides can be recovered from the host cells and rapidly sequenced using modern equipment. These methods allow the rapid determination of the importance of individual amino acid residues in a

polypeptide of interest, and can be applied to polypeptides of unknown structure.

Using the methods discussed above, one of ordinary skill in the art can identify and/or prepare a variety of polypeptides that are substantially homologous to residues 40 to 559 of SEQ ID NO: 2, 4 or 6 and retain the xyloglucanase activity of the wild-type protein.

The xyloglucanase enzyme of the invention may, in addition to the enzyme core comprising the catalytically domain, also comprise a cellulose binding domain (CBD), the cellulose binding domain and enzyme core (the catalytically active domain) of the enzyme being operably linked. The cellulose binding domain (CBD) may exist as an integral part the encoded enzyme, or a CBD from another origin may be introduced into the xyloglucanase thus creating an enzyme hybride. In this context, the term "cellulose-binding domain" is intended to be understood as defined by Peter Tomme et al. "Cellulose-Binding Domains: Classification and Properties" in "Enzymatic Degradation of Insoluble Carbohydrates", John N. Saddler and Michael H. Penner (Eds.), ACS Symposium Series, No. 618, 1996. This definition classifies more than 120 cellulose-binding domains into 10 families (I-X), and demonstrates that CBDs are found in various enzymes such as cellulases, xylanases, mannanases, arabinofuranosidases, acetyl esterases and chitinases. CBDs have also been found in algae, e.g. the red alga *Porphyra purpurea* as a non-hydrolytic polysaccharide-binding protein, see Tomme et al., op.cit. However, most of the CBDs are from cellulases and xylanases, CBDs are found at the N and C termini of proteins or are internal. Enzyme hybrids are known in the art, see e.g. WO 90/00609 and WO 95/16782, and may be prepared by transforming into a host cell a DNA construct comprising at least a fragment of DNA encoding the cellulose-binding domain ligated, with or without a linker, to a DNA sequence encoding the xyloglucanase and growing the host cell to express the fused gene. Enzyme hybrids may be described by the following formula:

CBD - MR - X

wherein CBD is the N-terminal or the C-terminal region of an amino acid sequence corresponding to at least the cellulose-binding domain; MR is the middle region (the linker), and may be

a bond, or a short linking group preferably of from about 2 to about 100 carbon atoms, more preferably of from 2 to 40 carbon atoms; or is preferably from about 2 to to about 100 amino acids, more preferably of from 2 to 40 amino acids; and X is an
5 N-terminal or C-terminal region of a polypeptide encoded by the polynucleotide molecule of the invention.

IMMUNOLOGICAL CROSS-REACTIVITY

Polyclonal antibodies, especially monospecific polyclonal
10 antibodies, to be used in determining immunological cross-reactivity may be prepared by use of a purified xyloglucanolytic enzyme. More specifically, antiserum against the xyloglucanase of the invention may be raised by immunizing rabbits (or other rodents) according to the procedure described by N. Axelsen et
15 al. in: A Manual of Quantitative Immunoelectrophoresis, Blackwell Scientific Publications, 1973, Chapter 23, or A. Johnstone and R. Thorpe, Immunochemistry in Practice, Blackwell Scientific Publications, 1982 (more specifically p. 27-31). Purified immunoglobulins may be obtained from the antisera, for
20 example by salt precipitation $((\text{NH}_4)_2\text{SO}_4)$, followed by dialysis and ion exchange chromatography, e.g. on DEAE-Sephadex. Immunochemical characterization of proteins may be done either by Outcherlony double-diffusion analysis (O. Ouchterlony in: Handbook of Experimental Immunology (D.M. Weir, Ed.), Blackwell
25 Scientific Publications, 1967, pp. 655-706), by crossed immunoelectrophoresis (N. Axelsen et al., supra, Chapters 3 and 4), or by rocket immunoelectrophoresis (N. Axelsen et al., Chapter 2).

30 RECOMBINANT EXPRESSION VECTORS

A recombinant vector comprising a DNA construct encoding the enzyme of the invention may be any vector which may conveniently be subjected to recombinant DNA procedures, and the choice of vector will often depend on the host cell into which it is to
35 be introduced. Thus, the vector may be an autonomously replicating vector, i.e. a vector which exists as an extrachromosomal entity, the replication of which is independent of chromosomal replication, e.g. a plasmid. Alternatively, the vector may be one which, when introduced into a host cell, is

integrated into the host cell genome in part or in its entirety and replicated together with the chromosome(s) into which it has been integrated.

The vector is preferably an expression vector in which the DNA sequence encoding the enzyme of the invention is operably linked to additional segments required for transcription of the DNA. In general, the expression vector is derived from plasmid or viral DNA, or may contain elements of both. The term, "operably linked" indicates that the segments are arranged so that they function in concert for their intended purposes, e.g. transcription initiates in a promoter and proceeds through the DNA sequence coding for the enzyme.

The promoter may be any DNA sequence which shows transcriptional activity in the host cell of choice and may be derived from genes encoding proteins either homologous or heterologous to the host cell.

Examples of suitable promoters for use in bacterial host cells include the promoter of the *Bacillus stearothermophilus* maltogenic amylase gene, the *Bacillus licheniformis* alpha-amylase gene, the *Bacillus amyloliquefaciens* alpha-amylase gene, the *Bacillus subtilis* alkaline protease gene, or the *Bacillus pumilus* xylosidase gene, or the phage Lambda P_R or P_L promoters or the *E. coli* lac, trp or tac promoters.

The DNA sequence encoding the enzyme of the invention may also, if necessary, be operably connected to a suitable terminator.

The recombinant vector of the invention may further comprise a DNA sequence enabling the vector to replicate in the host cell in question.

The vector may also comprise a selectable marker, e.g. a gene the product of which complements a defect in the host cell, or a gene encoding resistance to e.g. antibiotics like kanamycin, chloramphenicol, erythromycin, tetracycline, spectinomycin, or the like, or resistance to heavy metals or herbicides.

To direct an enzyme of the present invention into the secretory pathway of the host cells, a secretory signal sequence (also known as a leader sequence, prepro sequence or pre sequence) may be provided in the recombinant vector. The

secretory signal sequence is joined to the DNA sequence encoding the enzyme in the correct reading frame. Secretory signal sequences are commonly positioned 5' to the DNA sequence encoding the enzyme. The secretory signal sequence may be that
5 normally associated with the enzyme or may be from a gene encoding another secreted protein.

The procedures used to ligate the DNA sequences coding for the present enzyme, the promoter and optionally the terminator and/or secretory signal sequence, respectively, or to assemble
10 these sequences by suitable PCR amplification schemes, and to insert them into suitable vectors containing the information necessary for replication or integration, are well known to persons skilled in the art (cf., for instance, Sambrook et al., op.cit.).

15

HOST CELLS

The cloned DNA molecule introduced into the host cell may be either homologous or heterologous to the host in question. If homologous to the host cell, i.e. produced by the host cell in
20 nature, it will typically be operably connected to another promoter sequence or, if applicable, another secretory signal sequence and/or terminator sequence than in its natural environment. The term "homologous" is intended to include a DNA sequence encoding an enzyme native to the host organism in
25 question. The term "heterologous" is intended to include a DNA sequence not expressed by the host cell in nature. Thus, the DNA sequence may be from another organism, or it may be a synthetic sequence.

The host cell into which the cloned DNA molecule or the
30 recombinant vector of the invention is introduced may be any cell which is capable of producing the desired enzyme and includes bacteria, yeast, fungi and higher eukaryotic cells.

Examples of bacterial host cells which on cultivation are capable of producing the enzyme of the invention may be a
35 gram-positive bacteria such as a strain of *Bacillus*, in particular *Bacillus alkalophilus*, *Bacillus amyloliquefaciens*, *Bacillus brevis*, *Bacillus lautus*, *Bacillus lentus*, *Bacillus licheniformis*, *Bacillus circulans*, *Bacillus coagulans*, *Bacillus megatherium*, *Bacillus stearothermophilus*, *Bacillus subtilis* and

Bacillus thuringiensis, a strain of *Lactobacillus*, a strain of *Streptococcus*, a strain of *Streptomyces*, in particular *Streptomyces lividans* and *Streptomyces murinus*, or the host cell may be a gram-negative bacteria such as a strain of *Escherichia coli*.

The transformation of the bacteria may be effected by protoplast transformation, electroporation, conjugation, or by using competent cells in a manner known per se (cf. e.g. Sambrook et al., supra).

When expressing the enzyme in a bacteria such as *Escherichia coli*, the enzyme may be retained in the cytoplasm, typically as insoluble granules (known as inclusion bodies), or may be directed to the periplasmic space by a bacterial secretion sequence. In the former case, the cells are lysed and the granules are recovered and denatured after which the enzyme is refolded by diluting the denaturing agent. In the latter case, the enzyme may be recovered from the periplasmic space by disrupting the cells, e.g. by sonication or osmotic shock, to release the contents of the periplasmic space and recovering the enzyme.

When expressing the enzyme in a gram-positive bacteria such as a strain of *Bacillus* or a strain of *Streptomyces*, the enzyme may be retained in the cytoplasm, or may be directed to the extracellular medium by a bacterial secretion sequence.

Examples of a fungal host cell which on cultivation are capable of producing the enzyme of the invention is e.g. a strain of *Aspergillus* or *Fusarium*, in particular *Aspergillus awamori*, *Aspergillus nidulans*, *Aspergillus niger*, *Aspergillus oryzae*, and *Fusarium oxysporum*, and a strain of *Trichoderma*, preferably *Trichoderma harzianum*, *Trichoderma reesei* and *Trichoderma viride*.

Fungal cells may be transformed by a process involving protoplast formation and transformation of the protoplasts followed by regeneration of the cell wall in a manner known per se. The use of a strain of *Aspergillus* as a host cell is described in EP 238 023 (Novo Nordisk A/S), the contents of which are hereby incorporated by reference.

Examples of a host cell of yeast origin which on cultivation are capable of producing the enzyme of the invention

is e.g. a strain of *Hansenula* sp., a strain of *Kluyveromyces* sp., in particular *Kluyveromyces lactis* and *Kluyveromyces marcianus*, a strain of *Pichia* sp., a strain of *Saccharomyces*, in particular *Saccharomyces carlsbergensis*, *Saccharomyces* cerevisiae, *Saccharomyces kluyveri* and *Saccharomyces uvarum*, a strain of *Schizosaccharomyces* sp., in particular *Schizosaccharomyces pombe*, and a strain of *Yarrowia* sp., in particular *Yarrowia lipolytica*.

Examples of a host cell of plant origin which on cultivation are capable of producing the enzyme of the invention is e.g. a plant cell of *Solanum tuberosum* or *Nicotiana tabacum*.

METHOD OF PRODUCING A XYLOGLUCANOLYTIC ENZYME

In another aspect, the present invention also relates to a method of producing the enzyme preparation of the invention, the method comprising culturing a microorganism capable of producing the xyloglucanase under conditions permitting the production of the enzyme, and recovering the enzyme from the culture. Culturing may be carried out using conventional fermentation techniques, e.g. culturing in shake flasks or fermentors with agitation to ensure sufficient aeration on a growth medium inducing production of the xyloglucanase enzyme. The growth medium may contain a conventional N-source such as peptone, yeast extract or casamino acids, a reduced amount of a conventional C-source such as dextrose or sucrose, and an inducer such as xyloglucan or composit plant substrates such as cereal brans (e.g. wheat bran or rice husk). The recovery may be carried out using conventional techniques, e.g. separation of bio-mass and supernatant by centrifugation or filtration, recovery of the supernatant or disruption of cells if the enzyme of interest is intracellular, perhaps followed by further purification as described in EP 0 406 314 or by crystallization as described in WO 97/15660.

Further, the present invention provides a method of producing an isolated enzyme according to the invention, wherein a suitable host cell, which has been transformed with a DNA sequence encoding the enzyme, is cultured under conditions permitting the production of the enzyme, and the resulting enzyme is recovered from the culture.

As defined herein, an isolated polypeptide (e.g. an enzyme) is a polypeptide which is essentially free of other polypeptides, e.g., at least about 20% pure, preferably at least about 40% pure, more preferably about 60% pure, even more preferably about 80% pure, most preferably about 90% pure, and even most preferably about 95% pure, as determined by SDS-PAGE.

The term "isolated polypeptide" may alternatively be termed "purified polypeptide".

When an expression vector comprising a DNA sequence encoding the enzyme is transformed into a heterologous host cell it is possible to enable heterologous recombinant production of the enzyme of the invention.

Thereby it is possible to make a highly purified or monocomponent xyloglucanolytic composition, characterized in being free from homologous impurities.

In this context homologous impurities means any impurities (e.g. other polypeptides than the enzyme of the invention) which originate from the homologous cell where the enzyme of the invention is originally obtained from.

In the present invention the homologous host cell may be a strain of *Paenibacillus sp.* or *Paenibacillus polymyxa*.

The medium used to culture the transformed host cells may be any conventional medium suitable for growing the host cells in question. The expressed xyloglucanolytic enzyme may conveniently be secreted into the culture medium and may be recovered therefrom by well-known procedures including separating the cells from the medium by centrifugation or filtration, precipitating proteinaceous components of the medium by means of a salt such as ammonium sulphate, followed by chromatographic procedures such as ion exchange chromatography, affinity chromatography, or the like.

The present invention also relates to a transgenic plant, plant part or plant cell which has been transformed with a DNA sequence encoding the xyloglucanase of the invention so as to express and produce this enzyme in recoverable quantities. The enzyme may be recovered from the plant or plant part.

The transgenic plant can be dicotyledonous or monocotyledonous, for short a dicot or a monocot. Examples of monocot plants are grasses, such as meadow grass (blue grass,

Poa), forage grass such as festuca, lolium, temperate grass, such as Agrostis, and cereals, e.g. wheat, oats, rye, barley, rice, sorghum and maize (corn).

Examples of dicot plants are tobacco, legumes, such as lupins, potato, sugar beet, pea, bean and soybean, and cruciferous (family Brassicaceae), such as cauliflower, oil seed rape and the closely related model organism Arabidopsis thaliana.

Examples of plant parts are stem, callus, leaves, root, fruits, seeds, and tubers. In the present context, also specific plant tissues, such as chloroplast, apoplast, mitochondria, vacuole, peroxisomes and cytoplasm are considered to be a plant part. Furthermore, any plant cell, whatever the tissue origin, is considered to be a plant part.

Also included within the scope of the invention are the progeny of such plants, plant parts and plant cells.

The transgenic plant or plant cell expressing the enzyme of the invention may be constructed in accordance with methods known in the art. In short the plant or plant cell is constructed by incorporating one or more expression constructs encoding the enzyme of the invention into the plant host genome and propagating the resulting modified plant or plant cell into a transgenic plant or plant cell.

Conveniently, the expression construct is a DNA construct which comprises a gene encoding the enzyme of the invention in operable association with appropriate regulatory sequences required for expression of the gene in the plant or plant part of choice. Furthermore, the expression construct may comprise a selectable marker useful for identifying host cells into which the expression construct has been integrated and DNA sequences necessary for introduction of the construct into the plant in question (the latter depends on the DNA introduction method to be used).

The choice of regulatory sequences, such as promoter and terminator sequences and optionally signal or transit sequences is determined, e.g. based on when, where and how the enzyme is desired to be expressed. For instance, the expression of the gene encoding the enzyme of the invention may be constitutive or inducible, or may be developmental, stage or tissue specific,

and the gene product may be targeted to a specific tissue or plant part such as seeds or leaves. Regulatory sequences are eg described by Tague et al, Plant, Phys., 86, 506, 1988.

For constitutive expression the 35S-CaMV promoter may be
5 used (Franck et al., 1980. Cell 21: 285-294). Organ-specific promoters may eg be a promoter from storage sink tissues such as seeds, potato tubers, and fruits (Edwards & Coruzzi, 1990. Annu. Rev. Genet. 24: 275-303), or from metabolic sink tissues such as meristems (Ito et al., 1994. Plant Mol. Biol. 24: 863-878), a
10 seed specific promoter such as the glutelin, prolamin, globulin or albumin promoter from rice (Wu et al., Plant and Cell Physiology Vol. 39, No. 8 pp. 885-889 (1998)), a *Vicia faba* promoter from the legumin B4 and the unknown seed protein gene from *Vicia faba* described by Conrad U. et al, Journal of Plant
15 Physiology Vol. 152, No. 6 pp. 708-711 (1998), a promoter from a seed oil body protein (Chen et al., Plant and cell physiology vol. 39, No. 9 pp. 935-941 (1998), the storage protein napA promoter from Brassica napus, or any other seed specific promoter known in the art, eg as described in WO 91/14772.
20 Furthermore, the promoter may be a leaf specific promoter such as the rbcs promoter from rice or tomato (Kyoizuka et al., Plant Physiology Vol. 102, No. 3 pp. 991-1000 (1993), the chlorella virus adenine methyltransferase gene promoter (Mitra, A. and Higgins, DW, Plant Molecular Biology Vol. 26, No. 1 pp. 85-93
25 (1994), or the aldP gene promoter from rice (Kagaya et al., Molecular and General Genetics Vol. 248, No. 6 pp. 668-674 (1995), or a wound inducible promoter such as the potato pin2 promoter (Xu et al, Plant Molecular Biology Vol. 22, No. 4 pp. 573-588 (1993).

30 A promoter enhancer element may be used to achieve higher expression of the enzyme in the plant. For instance, the promoter enhancer element may be an intron placed between the promoter and the nucleotide sequence encoding the enzyme. For instance, Xu et al. op cit disclose the use of the first intron
35 of the rice actin 1 gene to enhance expression.

The selectable marker gene and any other parts of the expression construct may be chosen from those available in the art.

The DNA construct is incorporated into the plant genome

according to conventional techniques known in the art, including *Agrobacterium*-mediated transformation, virus-mediated transformation, micro injection, particle bombardment, biolistic transformation, and electroporation (Gasser et al, Science, 244, 5 1293; Potrykus, Bio/Techn. 8, 535, 1990; Shimamoto et al, Nature, 338, 274, 1989).

Presently, *Agrobacterium tumefaciens* mediated gene transfer is the method of choice for generating transgenic dicots (for review Hooykas & Schilperoort, 1992. Plant Mol. 10 Biol. 19: 15-38), however it can also be used for transforming monocots, although other transformation methods are generally preferred for these plants. Presently, the method of choice for generating transgenic monocots is particle bombardment (microscopic gold or tungsten particles coated with the 15 transforming DNA) of embryonic calli or developing embryos (Christou, 1992. Plant J. 2: 275-281; Shimamoto, 1994. Curr. Opin. Biotechnol. 5: 158-162; Vasil et al., 1992. Bio/Technology 10: 667-674). An alternative method for transformation of monocots is based on protoplast transformation as described by 20 Omirulleh S, et al., Plant Molecular biology Vol. 21, No. 3 pp. 415-428 (1993).

Following transformation, the transformants having incorporated the expression construct are selected and regenerated into whole plants according to methods well-known in 25 the art.

THE ENZYME

In a preferred embodiment of the present invention, the xyloglucanase has a relative activity at a temperature of 50°C, 30 preferably of at least 60%, preferably at least 70%, compared to the activity at the optimal temperature.

In yet another preferred embodiment, at a temperature of 60°C, the relative xyloglucanase activity is at least 40%, preferably at least 50%; at a temperature of 70°C, the relative 35 xyloglucanase activity is at least 40%, preferably at least 45%, especially at least 50%.

ENZYME COMPOSITIONS

In a still further aspect, the present invention relates to an enzyme composition comprising an enzyme exhibiting xyloglucanase activity as described above.

The enzyme composition of the invention may, in addition to the xyloglucanase of the invention, comprise one or more other enzyme types, for instance hemicellulase such as xylanase and mannanase, cellulase or endo- β -1,4-glucanase components, chitinase, lipase, esterase, pectinase, cutinase, phytase, oxidoreductase (peroxidase, haloperoxidase, oxidase, laccase), protease, amylase, reductase, phenoloxidase, ligninase, pullulanase, pectate lyase, pectin acetyl esterase, polygalacturonase, rhamnogalacturonase, pectin lyase, pectin methylesterase, cellobiohydrolase, transglutaminase; or mixtures thereof.

The enzyme composition may be prepared in accordance with methods known in the art and may be in the form of a liquid or a dry composition. For instance, the enzyme composition may be in the form of a granulate or a microgranulate. The enzyme to be included in the composition may be stabilized in accordance with methods known in the art.

Xyloglucanases have potential uses in a lot of different industries and applications. Examples are given below of preferred uses of the enzyme composition of the invention. The dosage of the enzyme composition of the invention and other conditions under which the composition is used may be determined based on methods known in the art.

The xyloglucanase or xyloglucanase composition according to the invention may be useful for at least one of the following purposes.

30

USES

Use in the detergent industry

During washing and wearing, dyestuff from dyed fabrics or garment will conventionally bleed from the fabric which then looks faded and worn. Removal of surface fibers from the fabric will partly restore the original colours and looks of the fabric. By the term "colour clarification", as used herein, is meant the partly restoration of the initial colours of fabric or garment throughout multiple washing cycles.

The term "de-pilling" denotes removing of pills from the fabric surface.

The term "soaking liquor" denotes an aqueous liquor in which laundry may be immersed prior to being subjected to a conventional washing process. The soaking liquor may contain one or more ingredients conventionally used in a washing or laundering process.

The term "washing liquor" denotes an aqueous liquor in which laundry is subjected to a washing process, i.e. usually a combined chemical and mechanical action either manually or in a washing machine. Conventionally, the washing liquor is an aqueous solution of a powder or liquid detergent composition.

The term "rinsing liquor" denotes an aqueous liquor in which laundry is immersed and treated, conventionally immediately after being subjected to a washing process, in order to rinse the laundry, i.e. essentially remove the detergent solution from the laundry. The rinsing liquor may contain a fabric conditioning or softening composition.

The laundry subjected to the method of the present invention may be conventional washable laundry. Preferably, the major part of the laundry is sewn or unsewn fabrics, including knits, wovens, denims, yarns, and toweling, made from cotton, cotton blends or natural or manmade cellulose (e.g. originating from xylan-containing cellulose fibers such as from wood pulp) or blends thereof. Examples of blends are blends of cotton or rayon/viscose with one or more companion material such as wool, synthetic fibers (e.g. polyamide fibers, acrylic fibers, polyester fibers, polyvinyl alcohol fibers, polyvinyl chloride fibers, polyvinylidene chloride fibers, polyurethane fibers, polyurea fibers, aramid fibers), and cellulose-containing fibers (e.g. rayon/viscose, ramie, flax/linen, jute, cellulose acetate fibers, lyocell).

DETERGENT DISCLOSURE AND EXAMPLES

35 Surfactant system

The detergent compositions according to the present invention comprise a surfactant system, wherein the surfactant can be selected from nonionic and/or anionic and/or cationic and/or ampholytic and/or zwitterionic and/or semi-polar

surfactants.

The surfactant is typically present at a level from 0.1% to 60% by weight.

The surfactant is preferably formulated to be compatible with enzyme components present in the composition. In liquid or gel compositions the surfactant is most preferably formulated in such a way that it promotes, or at least does not degrade, the stability of any enzyme in these compositions.

Preferred systems to be used according to the present invention comprise as a surfactant one or more of the nonionic and/or anionic surfactants described herein.


Polyethylene, polypropylene, and polybutylene oxide condensates of alkyl phenols are suitable for use as the nonionic surfactant of the surfactant systems of the present invention, with the polyethylene oxide condensates being preferred. These compounds include the condensation products of alkyl phenols having an alkyl group containing from about 6 to about 14 carbon atoms, preferably from about 8 to about 14 carbon atoms, in either a straight chain or branched-chain configuration with the alkylene oxide. In a preferred embodiment, the ethylene oxide is present in an amount equal to from about 2 to about 25 moles, more preferably from about 3 to about 15 moles, of ethylene oxide per mole of alkyl phenol. Commercially available nonionic surfactants of this type include Igepal™ CO-630, marketed by the GAF Corporation; and Triton™ X-45, X-114, X-100 and X-102, all marketed by the Rohm & Haas Company. These surfactants are commonly referred to as alkylphenol alkoxylates (e.g., alkyl phenol ethoxylates).

The condensation products of primary and secondary aliphatic alcohols with about 1 to about 25 moles of ethylene oxide are suitable for use as the nonionic surfactant of the nonionic surfactant systems of the present invention. The alkyl chain of the aliphatic alcohol can either be straight or branched, primary or secondary, and generally contains from about 8 to about 22 carbon atoms. Preferred are the condensation products of alcohols having an alkyl group containing from about 8 to about 20 carbon atoms, more preferably from about 10 to about 18 carbon atoms, with from about 2 to about 10 moles of ethylene oxide per mole of alcohol. About 2 to about 7 moles of

ethylene oxide and most preferably from 2 to 5 moles of ethylene oxide per mole of alcohol are present in said condensation products. Examples of commercially available nonionic surfactants of this type include Tergitol™ 15-S-9 (The condensation product of C₁₁-C₁₅ linear alcohol with 9 moles ethylene oxide), Tergitol™ 24-L-6 NMW (the condensation product of C₁₂-C₁₄ primary alcohol with 6 moles ethylene oxide with a narrow molecular weight distribution), both marketed by Union Carbide Corporation; Neodol™ 45-9 (the condensation product of C₁₄-C₁₅ linear alcohol with 9 moles of ethylene oxide), Neodol™ 23-3 (the condensation product of C₁₂-C₁₃ linear alcohol with 3.0 moles of ethylene oxide), Neodol™ 45-7 (the condensation product of C₁₄-C₁₅ linear alcohol with 7 moles of ethylene oxide), Neodol™ 45-5 (the condensation product of C₁₄-C₁₅ linear alcohol with 5 moles of ethylene oxide) marketed by Shell Chemical Company, Kyro™ EOB (the condensation product of C₁₃-C₁₅ alcohol with 9 moles ethylene oxide), marketed by The Procter & Gamble Company, and Genapol LA 050 (the condensation product of C₁₂-C₁₄ alcohol with 5 moles of ethylene oxide) marketed by Hoechst. Preferred range of HLB in these products is from 8-11 and most preferred from 8-10.

Also useful as the nonionic surfactant of the surfactant systems of the present invention are alkylpolysaccharides disclosed in US 4,565,647, having a hydrophobic group containing from about 6 to about 30 carbon atoms, preferably from about 10 to about 16 carbon atoms and a polysaccharide, e.g. a polyglycoside, hydrophilic group containing from about 1.3 to about 10, preferably from about 1.3 to about 3, most preferably from about 1.3 to about 2.7 saccharide units. Any reducing saccharide containing 5 or 6 carbon atoms can be used, e.g., glucose, galactose and galactosyl moieties can be substituted for the glucosyl moieties (optionally the hydrophobic group is attached at the 2-, 3-, 4-, etc. positions thus giving a glucose or galactose as opposed to a glucoside or galactoside). The intersaccharide bonds can be, e.g., between the one position of the additional saccharide units and the 2-, 3-, 4-, and/or 6-positions on the preceding saccharide units.

The preferred alkylpolyglycosides have the formula




wherein R^2 is selected from the group consisting of alkyl, alkylphenyl, hydroxyalkyl, hydroxyalkylphenyl, and mixtures thereof in which the alkyl groups contain from about 10 to about 18, preferably from about 12 to about 14, carbon atoms; n is 2 or 3, preferably 2; t is from 0 to about 10, preferably 0; and x is from about 1.3 to about 10, preferably from about 1.3 to about 3, most preferably from about 1.3 to about 2.7. The glycosyl is preferably derived from glucose. To prepare these compounds, the alcohol or alkylpolyethoxy alcohol is formed first and then reacted with glucose, or a source of glucose, to form the glucoside (attachment at the 1-position). The additional glycosyl units can then be attached between their 1-position and the preceding glycosyl units 2-, 3-, 4-, and/or 6-position, preferably predominantly the 2-position.

The condensation products of ethylene oxide with a hydrophobic base formed by the condensation of propylene oxide with propylene glycol are also suitable for use as the additional nonionic surfactant systems of the present invention. The hydrophobic portion of these compounds will preferably have a molecular weight from about 1500 to about 1800 and will exhibit water insolubility. The addition of polyoxyethylene moieties to this hydrophobic portion tends to increase the water solubility of the molecule as a whole, and the liquid character of the product is retained up to the point where the polyoxyethylene content is about 50% of the total weight of the condensation product, which corresponds to condensation with up to about 40 moles of ethylene oxide. Examples of compounds of this type include certain of the commercially available Pluronic™ surfactants, marketed by BASF.

Also suitable for use as the nonionic surfactant of the nonionic surfactant system of the present invention, are the condensation products of ethylene oxide with the product resulting from the reaction of propylene oxide and ethylenediamine. The hydrophobic moiety of these products consists of the reaction product of ethylenediamine and excess propylene oxide, and generally has a molecular weight of from about 2500 to about 3000. This hydrophobic moiety is condensed

with ethylene oxide to the extent that the condensation product contains from about 40% to about 80% by weight of polyoxyethylene and has a molecular weight of from about 5,000 to about 11,000. Examples of this type of nonionic surfactant include certain of the commercially available Tetronic™ compounds, marketed by BASF.

Preferred for use as the nonionic surfactant of the surfactant systems of the present invention are polyethylene oxide condensates of alkyl phenols, condensation products of primary and secondary aliphatic alcohols with from about 1 to about 25 moles of ethyleneoxide, alkylpolysaccharides, and mixtures hereof. Most preferred are C₈-C₁₄ alkyl phenol ethoxylates having from 3 to 15 ethoxy groups and C₈-C₁₈ alcohol ethoxylates (preferably C₁₀ avg.) having from 2 to 10 ethoxy groups, and mixtures thereof. Highly preferred nonionic surfactants are polyhydroxy fatty acid amide surfactants of the formula



wherein R¹ is H, or R¹ is C₁₋₄ hydrocarbyl, 2-hydroxyethyl, 2-hydroxypropyl or a mixture thereof, R² is C₅₋₃₁ hydrocarbyl, and Z is a polyhydroxyhydrocarbyl having a linear hydrocarbyl chain with at least 3 hydroxyls directly connected to the chain, or an alkoxyated derivative thereof. Preferably, R¹ is methyl, R² is straight C₁₁₋₁₅ alkyl or C₁₆₋₁₈ alkyl or alkenyl chain such as coconut alkyl or mixtures thereof, and Z is derived from a reducing sugar such as glucose, fructose, maltose or lactose, in a reductive amination reaction.

Highly preferred anionic surfactants include alkyl alkoxyated sulfate surfactants. Examples hereof are water soluble salts or acids of the formula RO(A)_mSO₃M wherein R is an unsubstituted C₁₀-C₂₄ alkyl or hydroxyalkyl group having a C₁₀-C₂₄ alkyl component, preferably a C₁₂-C₂₀ alkyl or hydro-xyalkyl, more preferably C₁₂-C₁₈ alkyl or hydroxyalkyl, A is an ethoxy or propoxy unit, m is greater than zero, typically between about 0.5 and about 6, more preferably between about 0.5 and about 3, and M is H or a cation which can be, for example, a metal cation (e.g., sodium, potassium, lithium, calcium, magnesium, etc.),

ammonium or substituted-ammonium cation. Alkyl ethoxylated sulfates as well as alkyl propoxylated sulfates are contemplated herein. Specific examples of substituted ammonium cations include methyl-, dimethyl, trimethyl-ammonium cations and

5 quaternary ammonium cations such as tetramethyl-ammonium and dimethyl piperdinium cations and those derived from alkylamines such as ethylamine, diethylamine, triethylamine, mixtures thereof, and the like. Exemplary surfactants are C_{12} - C_{18} alkyl polyethoxylate (1.0) sulfate (C_{12} - $C_{18}E(1.0)M$), C_{12} - C_{18} alkyl


10 polyethoxylate (2.25) sulfate (C_{12} - $C_{18}(2.25)M$, and C_{12} - C_{18} alkyl polyethoxylate (3.0) sulfate (C_{12} - $C_{18}E(3.0)M$), and C_{12} - C_{18} alkyl polyethoxylate (4.0) sulfate (C_{12} - $C_{18}E(4.0)M$), wherein M is conveniently selected from sodium and potassium.

Suitable anionic surfactants to be used are alkyl ester

15 sulfonate surfactants including linear esters of C_8 - C_{20} carboxylic acids (i.e., fatty acids) which are sulfonated with gaseous SO_3 according to "The Journal of the American Oil Chemists Society", 52 (1975), pp. 323-329. Suitable starting materials would include natural fatty substances as derived from

20 tallow, palm oil, etc.

The preferred alkyl ester sulfonate surfactant, especially for laundry applications, comprise alkyl ester sulfonate surfactants of the structural formula:



30 wherein R^3 is a C_8 - C_{20} hydrocarbyl, preferably an alkyl, or combination thereof, R^4 is a C_1 - C_6 hydrocarbyl, preferably an alkyl, or combination thereof, and M is a cation which forms a water soluble salt with the alkyl ester sulfonate. Suitable salt-forming cations include metals such as sodium, potassium,

35 and lithium, and substituted or unsubstituted ammonium cations, such as monoethanolamine, diethanolamine, and triethanolamine. Preferably, R^3 is C_{10} - C_{16} alkyl, and R^4 is methyl, ethyl or isopropyl. Especially preferred are the methyl ester sulfonates wherein R^3 is C_{10} - C_{16} alkyl.

Other suitable anionic surfactants include the alkyl sulfate surfactants which are water soluble salts or acids of the formula ROSO_3M wherein R preferably is a $\text{C}_{10}\text{-C}_{24}$ hydrocarbyl, preferably an alkyl or hydroxyalkyl having a $\text{C}_{10}\text{-C}_{20}$ alkyl component, more preferably a $\text{C}_{12}\text{-C}_{18}$ alkyl or hydroxyalkyl, and M is H or a cation, e.g., an alkali metal cation (e.g. sodium, potassium, lithium), or ammonium or substituted ammonium (e.g. methyl-, dimethyl-, and trimethyl ammonium cations and quaternary ammonium cations such as tetramethyl-ammonium and dimethyl piperdinium cations and quaternary ammonium cations derived from alkylamines such as ethylamine, diethylamine, triethylamine, and mixtures thereof, and the like). Typically, alkyl chains of $\text{C}_{12}\text{-C}_{16}$ are preferred for lower wash temperatures (e.g. below about 50°C) and $\text{C}_{16}\text{-C}_{18}$ alkyl chains are preferred for higher wash temperatures (e.g. above about 50°C).

Other anionic surfactants useful for deterative purposes can also be included in the laundry detergent compositions of the present invention. These can include salts (including, for example, sodium, potassium, ammonium, and substituted ammonium salts such as mono- di- and triethanolamine salts) of soap, $\text{C}_8\text{-C}_{22}$ primary or secondary alkanesulfonates, $\text{C}_8\text{-C}_{24}$ olefinsulfonates, sulfonated polycarboxylic acids prepared by sulfonation of the pyrolyzed product of alkaline earth metal citrates, e.g., as described in British patent specification No. 1,082,179, $\text{C}_8\text{-C}_{24}$ alkylpolyglycolethersulfates (containing up to 10 moles of ethylene oxide); alkyl glycerol sulfonates, fatty acyl glycerol sulfonates, fatty oleyl glycerol sulfates, alkyl phenol ethylene oxide ether sulfates, paraffin sulfonates, alkyl phosphates, isethionates such as the acyl isethionates, N-acyl taurates, alkyl succinamates and sulfosuccinates, monoesters of sulfosuccinates (especially saturated and unsaturated $\text{C}_{12}\text{-C}_{18}$ monoesters) and diesters of sulfosuccinates (especially saturated and unsaturated $\text{C}_6\text{-C}_{12}$ diesters), acyl sarcosinates, sulfates of alkylpolysaccharides such as the sulfates of alkylpolyglucoside (the nonionic nonsulfated compounds being described below), branched primary alkyl sulfates, and alkyl polyethoxy carboxylates such as those of the formula $\text{RO}(\text{CH}_2\text{CH}_2\text{O})_k\text{-CH}_2\text{COO-M}^+$ wherein R is a $\text{C}_8\text{-C}_{22}$ alkyl, k is an integer from 1 to 10, and M is a soluble salt forming cation. Resin

acids and hydrogenated resin acids are also suitable, such as rosin, hydrogenated rosin, and resin acids and hydrogenated resin acids present in or derived from tall oil.


Alkylbenzene sulfonates are highly preferred. Especially
5 preferred are linear (straight-chain) alkyl benzene sulfonates (LAS) wherein the alkyl group preferably contains from 10 to 18 carbon atoms.

Further examples are described in "Surface Active Agents and Detergents" (Vol. I and II by Schwartz, Perry and Berch). A
10 variety of such surfactants are also generally disclosed in US 3,929,678, (Column 23, line 58 through Column 29, line 23, herein incorporated by reference).

When included therein, the laundry detergent compositions of the present invention typically comprise from about 1% to
15 about 40%, preferably from about 3% to about 20% by weight of such anionic surfactants.

The laundry detergent compositions of the present invention may also contain cationic, ampholytic, zwitterionic, and semi-polar surfactants, as well as the nonionic and/or
20 anionic surfactants other than those already described herein.

Cationic deterative surfactants suitable for use in the laundry detergent compositions of the present invention are those having one long-chain hydrocarbyl group. Examples of such cationic surfactants include the ammonium surfactants such as
25 alkyltrimethylammonium halogenides, and those surfactants having the formula:



wherein R^2 is an alkyl or alkyl benzyl group having from about 8 to about 18 carbon atoms in the alkyl chain, each R^3 is selected from the group consisting of $-CH_2CH_2-$, $-CH_2CH(CH_3)-$, $-CH_2CH(CH_2OH)-$, $-CH_2CH_2CH_2-$, and mixtures thereof; each R^4 is selected from the group consisting of C_1-C_4 alkyl, C_1-C_4 hydroxyalkyl, benzyl ring structures formed by joining the two R^4 groups, $-CH_2CHOHCHOHCOR^6CHOHCH_2OH$, wherein R^6 is any hexose or hexose polymer having a molecular weight less than about 1000, and hydrogen when y is not 0; R^5 is the same as R^4 or is an alkyl chain, wherein the total number of carbon atoms or R^2 plus R^5 is not more than about 18; each y is from 0 to about 10, and the sum of the y values is from 0 to about 15; and X is any compatible anion.

Highly preferred cationic surfactants are the water soluble quaternary ammonium compounds useful in the present composition having the formula:



wherein R_1 is C_8-C_{16} alkyl, each of R_2 , R_3 and R_4 is independently C_1-C_4 alkyl, C_1-C_4 hydroxy alkyl, benzyl, and $-(C_2H_4)_xH$ where x has a value from 2 to 5, and X is an anion. Not more than one of R_2 , R_3 or R_4 should be benzyl.


The preferred alkyl chain length for R_1 is $C_{12}-C_{15}$, particularly where the alkyl group is a mixture of chain lengths derived from coconut or palm kernel fat or is derived synthetically by olefin build up or OXO alcohols synthesis.

Preferred groups for R_2 , R_3 and R_4 are methyl and hydroxyethyl groups and the anion X may be selected from halide, methosulphate, acetate and phosphate ions.

Examples of suitable quaternary ammonium compounds of formulae (i) for use herein are:

coconut trimethyl ammonium chloride or bromide;
 coconut methyl dihydroxyethyl ammonium chloride or bromide;
 decyl triethyl ammonium chloride;
 decyl dimethyl hydroxyethyl ammonium chloride or bromide;
 C_{12-15} dimethyl hydroxyethyl ammonium chloride or bromide;
 coconut dimethyl hydroxyethyl ammonium chloride or bromide;
 myristyl trimethyl ammonium methyl sulphate;

lauryl dimethyl benzyl ammonium chloride or bromide;
 lauryl dimethyl (ethenoxy), ammonium chloride or bromide;
 choline esters (compounds of formula (i) wherein R_1 is



di-alkyl imidazolines [compounds of formula (i)].

10 Other cationic surfactants useful herein are also described in US 4,228,044 and in EP 000 224.

When included therein, the laundry detergent compositions of the present invention typically comprise from 0.2% to about 25%, preferably from about 1% to about 8% by weight of such
 15 cationic surfactants.

Ampholytic surfactants are also suitable for use in the laundry detergent compositions of the present invention. These surfactants can be broadly described as aliphatic derivatives of secondary or tertiary amines, or aliphatic derivatives of
 20 heterocyclic secondary and tertiary amines in which the aliphatic radical can be straight- or branched-chain. One of the aliphatic substituents contains at least about 8 carbon atoms, typically from about 8 to about 18 carbon atoms, and at least one contains an anionic water-solubilizing group, e.g. carboxy,
 25 sulfonate, sulfate. See US 3,929,678 (column 19, lines 18-35) for examples of ampholytic surfactants.

When included therein, the laundry detergent compositions of the present invention typically comprise from 0.2% to about 15%, preferably from about 1% to about 10% by weight of such
 30 ampholytic surfactants.

Zwitterionic surfactants are also suitable for use in laundry detergent compositions. These surfactants can be broadly described as derivatives of secondary and tertiary amines, derivatives of heterocyclic secondary and tertiary amines, or
 35 derivatives of quaternary ammonium, quaternary phosphonium or tertiary sulfonium compounds. See US 3,929,678 (column 19, line 38 through column 22, line 48) for examples of zwitterionic surfactants.


When included therein, the laundry detergent compositions

of the present invention typically comprise from 0.2% to about 15%, preferably from about 1% to about 10% by weight of such zwitterionic surfactants.

Semi-polar nonionic surfactants are a special category of nonionic surfactants which include water-soluble amine oxides containing one alkyl moiety of from about 10 to about 18 carbon atoms and 2 moieties selected from the group consisting of alkyl groups and hydroxyalkyl groups containing from about 1 to about 3 carbon atoms; watersoluble phosphine oxides containing one alkyl moiety of from about 10 to about 18 carbon atoms and 2 moieties selected from the group consisting of alkyl groups and hydroxyalkyl groups containing from about 1 to about 3 carbon atoms; and water-soluble sulfoxides containing one alkyl moiety from about 10 to about 18 carbon atoms and a moiety selected from the group consisting of alkyl and hydroxyalkyl moieties of from about 1 to about 3 carbon atoms.

Semi-polar nonionic detergent surfactants include the amine oxide surfactants having the formula:

20



wherein R^3 is an alkyl, hydroxyalkyl, or alkyl phenyl group or mixtures thereof containing from about 8 to about 22 carbon atoms; R^4 is an alkylene or hydroxyalkylene group containing from about 2 to about 3 carbon atoms or mixtures thereof; x is from 0 to about 3; and each R^5 is an alkyl or hydroxyalkyl group containing from about 1 to about 3 carbon atoms or a polyethylene oxide group containing from about 1 to about 3 ethylene oxide groups. The R^5 groups can be attached to each other, e.g., through an oxygen or nitrogen atom, to form a ring structure.

These amine oxide surfactants in particular include C_{10} - C_{18} alkyl dimethyl amine oxides and C_8 - C_{12} alkoxy ethyl dihydroxy ethyl amine oxides.

When included therein, the laundry detergent compositions of the present invention typically comprise from 0.2% to about 15%, preferably from about 1% to about 10% by weight of such

semi-polar nonionic surfactants.

Builder system

The compositions according to the present invention may
5 further comprise a builder system. Any conventional builder
system is suitable for use herein including aluminosilicate
materials, silicates, polycarboxylates and fatty acids,
materials such as ethylenediamine tetraacetate, metal ion
sequestrants such as aminopolyphosphonates, particularly
10 ethylenediamine tetramethylene phosphonic acid and diethylene
triamine pentamethylenephosphonic acid. Though less preferred
for obvious environmental reasons, phosphate builders can also
be used herein.

Suitable builders can be an inorganic ion exchange
15 material, commonly an inorganic hydrated aluminosilicate
material, more particularly a hydrated synthetic zeolite such as
hydrated zeolite A, X, B, HS or MAP.

Another suitable inorganic builder material is layered
silicate, e.g. SKS-6 (Hoechst). SKS-6 is a crystalline layered
20 silicate consisting of sodium silicate ($\text{Na}_2\text{Si}_2\text{O}_5$).

Suitable polycarboxylates containing one carboxy group
include lactic acid, glycolic acid and ether derivatives thereof
as disclosed in Belgian Patent Nos. 831,368, 821,369 and
821,370. Polycarboxylates containing two carboxy groups include
25 the water-soluble salts of succinic acid, malonic acid,
(ethylenedioxy) diacetic acid, maleic acid, diglycollic acid,
tartaric acid, tartronic acid and fumaric acid, as well as the
ether carboxylates described in German Offenle-enschrift
2,446,686, and 2,446,487, US 3,935,257 and the sulfinyl
30 carboxylates described in Belgian Patent No. 840,623.

Polycarboxylates containing three carboxy groups include, in
particular, water-soluble citrates, aconitrates and citraconates
as well as succinate derivatives such as the
carboxymethyloxysuccinates described in British Patent No.
35 1,379,241, lactoxysuccinates described in Netherlands
Application 7205873, and the oxypolycarboxylate materials such
as 2-oxa-1,1,3-propane tricarboxylates described in British
Patent No. 1,387,447.

Polycarboxylates containing four carboxy groups include

oxydisuccinates disclosed in British Patent No. 1,261,829, 1,1,2,2,-ethane tetracarboxylates, 1,1,3,3-propane tetracarboxylates containing sulfo substituents include the sulfosuccinate derivatives disclosed in British Patent Nos.

5 1,398,421 and 1,398,422 and in US 3,936,448, and the sulfonated pyrolysed citrates described in British Patent No. 1,082,179, while polycarboxylates containing phosphone substituents are disclosed in British Patent No. 1,439,000.

Alicyclic and heterocyclic polycarboxylates include
10 cyclopentane-cis,cis-cis-tetracarboxylates, cyclopentadienide pentacarboxylates, 2,3,4,5-tetrahydro-furan - cis, cis, cis-tetracarboxylates, 2,5-tetrahydro-furan-cis, dicarboxylates, 2,2,5,5,-tetrahydrofuran - tetracarboxylates, 1,2,3,4,5,6-hexane - hexacarboxylates and carboxymethyl derivatives of polyhydric
15 alcohols such as sorbitol, mannitol and xylitol. Aromatic polycarboxylates include mellitic acid, pyromellitic acid and the phthalic acid derivatives disclosed in British Patent No. 1,425,343.

Of the above, the preferred polycarboxylates are hydroxy-
20 carboxylates containing up to three carboxy groups per molecule, more particularly citrates.

Preferred builder systems for use in the present compositions include a mixture of a water-insoluble aluminosilicate builder such as zeolite A or of a layered
25 silicate (SKS-6), and a water-soluble carboxylate chelating agent such as citric acid.

A suitable chelant for inclusion in the detergent compositions in accordance with the invention is ethylenediamine-N,N'-disuccinic acid (EDDS) or the alkali metal, alkaline earth
30 metal, ammonium, or substituted ammonium salts thereof, or mixtures thereof. Preferred EDDS compounds are the free acid form and the sodium or magnesium salt thereof. Examples of such preferred sodium salts of EDDS include Na₂EDDS and Na₄EDDS. Examples of such preferred magnesium salts of EDDS include
35 MgEDDS and Mg₂EDDS. The magnesium salts are the most preferred for inclusion in compositions in accordance with the invention.

Preferred builder systems include a mixture of a water-insoluble aluminosilicate builder such as zeolite A, and a water soluble carboxylate chelating agent such as citric acid.

Other builder materials that can form part of the builder system for use in granular compositions include inorganic materials such as alkali metal carbonates, bicarbonates, silicates, and organic materials such as the organic
5 phosphonates, amino polyalkylene phosphonates and amino polycarboxylates.

Other suitable water-soluble organic salts are the homo- or co-polymeric acids or their salts, in which the polycarboxylic acid comprises at least two carboxyl radicals
10 separated from each other by not more than two carbon atoms.

Polymers of this type are disclosed in GB-A-1,596,756. Examples of such salts are polyacrylates of MW 2000-5000 and their copolymers with maleic anhydride, such copolymers having a molecular weight of from 20,000 to 70,000, especially about
15 40,000.

Detergency builder salts are normally included in amounts of from 5% to 80% by weight of the composition. Preferred levels of builder for liquid detergents are from 5% to 30%.

20 Enzymes

Preferred detergent compositions, in addition to the enzyme preparation of the invention, comprise other enzyme(s) which provides cleaning performance and/or fabric care benefits.

Such enzymes include proteases, lipases, cutinases,
25 amylases, cellulases, peroxidases, oxidases (e.g. laccases).

Proteases: Any protease suitable for use in alkaline solutions can be used. Suitable proteases include those of animal, vegetable or microbial origin. Microbial origin is preferred. Chemically or genetically modified mutants are included. The
30 protease may be a serine protease, preferably an alkaline microbial protease or a trypsin-like protease. Examples of alkaline proteases are subtilisins, especially those derived from Bacillus, e.g., subtilisin Novo, subtilisin Carlsberg, subtilisin 309, subtilisin 147 and subtilisin 168 (described in
35 WO 89/06279). Examples of trypsin-like proteases are trypsin (e.g. of porcine or bovine origin) and the Fusarium protease described in WO 89/06270.

Preferred commercially available protease enzymes include those sold under the trade names Alcalase, Savinase, Primase,

Durazym, and Esperase by Novo Nordisk A/S (Denmark), those sold under the tradename Maxatase, Maxacal, Maxapem, Properase, Purafect and Purafect OXP by Genencor International, and those sold under the tradename Opticlean and Optimase by Solvay

5 Enzymes. Protease enzymes may be incorporated into the compositions in accordance with the invention at a level of from 0.00001% to 2% of enzyme protein by weight of the composition, preferably at a level of from 0.0001% to 1% of enzyme protein by weight of the composition, more preferably at a level of from
10 0.001% to 0.5% of enzyme protein by weight of the composition, even more preferably at a level of from 0.01% to 0.2% of enzyme protein by weight of the composition.

Lipases: Any lipase suitable for use in alkaline solutions can be used. Suitable lipases include those of bacterial or
15 fungal origin. Chemically or genetically modified mutants are included.

Examples of useful lipases include a Humicola lanuginosa lipase, e.g., as described in EP 258 068 and EP 305 216, a Rhizomucor miehei lipase, e.g., as described in EP 238 023, a
20 Candida lipase, such as a C. antarctica lipase, e.g., the C. antarctica lipase A or B described in EP 214 761, a Pseudomonas lipase such as a P. alcaligenes and P. pseudoalcaligenes lipase, e.g., as described in EP 218 272, a P. cepacia lipase, e.g., as described in EP 331 376, a P. stutzeri lipase, e.g., as
25 disclosed in GB 1,372,034, a P. fluorescens lipase, a Bacillus lipase, e.g., a B. subtilis lipase (Dartois et al., (1993), Biochemica et Biophysica acta 1131, 253-260), a B. stearo-thermophilus lipase (JP 64/744992) and a B. pumilus lipase (WO 91/16422).

30 Furthermore, a number of cloned lipases may be useful, including the Penicillium camembertii lipase described by Yamaguchi et al., (1991), Gene 103, 61-67), the Geotricum candidum lipase (Schimada, Y. et al., (1989), J. Biochem., 106, 383-388), and various Rhizopus lipases such as a R. delemar
35 lipase (Hass, M.J et al., (1991), Gene 109, 117-113), a R. niveus lipase (Kugimiya et al., (1992), Biosci. Biotech. Biochem. 56, 716-719) and a R. oryzae lipase.

Other types of lipolytic enzymes such as cutinases may also be useful, e.g., a cutinase derived from Pseudomonas

mendocina as described in WO 88/09367, or a cutinase derived from Fusarium solani pisi (e.g. described in WO 90/09446).

Especially suitable lipases are lipases such as M1 Lipase™, Luma fast™ and Lipomax™ (Genencor), Lipolase™ and
5 Lipolase Ultra™ (Novo Nordisk A/S), and Lipase P "Amano" (Amano Pharmaceutical Co. Ltd.).

The lipases are normally incorporated in the detergent composition at a level of from 0.00001% to 2% of enzyme protein by weight of the composition, preferably at a level of from
10 0.0001% to 1% of enzyme protein by weight of the composition, more preferably at a level of from 0.001% to 0.5% of enzyme protein by weight of the composition, even more preferably at a level of from 0.01% to 0.2% of enzyme protein by weight of the composition.

15 Amylases: Any amylase (a and/or b) suitable for use in alkaline solutions can be used. Suitable amylases include those of bacterial or fungal origin. Chemically or genetically modified mutants are included. Amylases include, for example, α -amylases obtained from a special strain of B. licheniformis,
20 described in more detail in GB 1,296,839. Commercially available amylases are Duramyl™, Termamyl™, Fungamyl™ and BAN™ (available from Novo Nordisk A/S) and Rapidase™ and Maxamyl P™ (available from Genencor).

The amylases are normally incorporated in the detergent
25 composition at a level of from 0.00001% to 2% of enzyme protein by weight of the composition, preferably at a level of from 0.0001% to 1% of enzyme protein by weight of the composition, more preferably at a level of from 0.001% to 0.5% of enzyme protein by weight of the composition, even more preferably at a
30 level of from 0.01% to 0.2% of enzyme protein by weight of the composition.

Cellulases: Any cellulase suitable for use in alkaline solutions can be used. Suitable cellulases include those of bacterial or fungal origin. Chemically or genetically modified
35 mutants are included. Suitable cellulases are disclosed in US 4,435,307 which discloses fungal cellulases produced from *Humicola insolens*, in WO 96/34108 and WO 96/34092 which disclose bacterial alkalophilic cellulases (BCE 103) from *Bacillus*, and in WO 94/21801, US 5,475,101 and US 5,419,778 which disclose EG

III cellulases from *Trichoderma*. Especially suitable cellulases are the cellulases having colour care benefits. Examples of such cellulases are cellulases described in European patent application No. 0 495 257. Commercially available cellulases include Celluzyme™ and Carezyme™ produced by a strain of *Humicola insolens* (Novo Nordisk A/S), KAC-500(B)™ (Kao Corporation), and Puradax™ (Genencor International).

Cellulases are normally incorporated in the detergent composition at a level of from 0.00001% to 2% of enzyme protein by weight of the composition, preferably at a level of from 0.0001% to 1% of enzyme protein by weight of the composition, more preferably at a level of from 0.001% to 0.5% of enzyme protein by weight of the composition, even more preferably at a level of from 0.01% to 0.2% of enzyme protein by weight of the composition.

Peroxidases/Oxidases : Peroxidase enzymes are used in combination with hydrogen peroxide or a source thereof (e.g. a percarbonate, perborate or persulfate). Oxidase enzymes are used in combination with oxygen. Both types of enzymes are used for "solution bleaching", i.e. to prevent transfer of a textile dye from a dyed fabric to another fabric when said fabrics are washed together in a wash liquor, preferably together with an enhancing agent as described in e.g. WO 94/12621 and WO 95/01426. Suitable peroxidases/oxidases include those of plant, bacterial or fungal origin. Chemically or genetically modified mutants are included.

Peroxidase and/or oxidase enzymes are normally incorporated in the detergent composition at a level of from 0.00001% to 2% of enzyme protein by weight of the composition, preferably at a level of from 0.0001% to 1% of enzyme protein by weight of the composition, more preferably at a level of from 0.001% to 0.5% of enzyme protein by weight of the composition, even more preferably at a level of from 0.01% to 0.2% of enzyme protein by weight of the composition.

Mixtures of the above mentioned enzymes are encompassed herein, in particular a mixture of a protease, an amylase, a lipase and/or a cellulase.

The enzyme of the invention, or any other enzyme incorporated in the detergent composition, is normally

incorporated in the detergent composition at a level from 0.00001% to 2% of enzyme protein by weight of the composition, preferably at a level from 0.0001% to 1% of enzyme protein by weight of the composition, more preferably at a level from 5 0.001% to 0.5% of enzyme protein by weight of the composition, even more preferably at a level from 0.01% to 0.2% of enzyme protein by weight of the composition.

Bleaching agents

10 Additional optional detergent ingredients that can be included in the detergent compositions of the present invention include bleaching agents such as PB1, PB4 and percarbonate with a particle size of 400-800 microns. These bleaching agent components can include one or more oxygen bleaching agents and, 15 depending upon the bleaching agent chosen, one or more bleach activators. When present oxygen bleaching compounds will typically be present at levels of from about 1% to about 25%. In general, bleaching compounds are optional added components in non-liquid formulations, e.g. granular detergents.

20 The bleaching agent component for use herein can be any of the bleaching agents useful for detergent compositions including oxygen bleaches as well as others known in the art.

The bleaching agent suitable for the present invention can be an activated or non-activated bleaching agent.

25 One category of oxygen bleaching agent that can be used encompasses percarboxylic acid bleaching agents and salts thereof. Suitable examples of this class of agents include magnesium monoperoxyphthalate hexahydrate, the magnesium salt of meta-chloro perbenzoic acid, 4-nonylamino-4-oxoperoxybutyric 30 acid and diperoxydodecanedioic acid. Such bleaching agents are disclosed in US 4,483,781, US 740,446, EP 0 133 354 and US 4,412,934. Highly preferred bleaching agents also include 6-nonylamino-6-oxoperoxy-caproic acid as described in US 4,634,551.

Another category of bleaching agents that can be used 35 encompasses the halogen bleaching agents. Examples of hypohalite bleaching agents, for example, include trichloro isocyanuric acid and the sodium and potassium dichloroisocyanurates and N-chloro and N-bromo alkane sulphonamides. Such materials are normally added at 0.5-10% by weight of the finished product,

preferably 1-5% by weight.

The hydrogen peroxide releasing agents can be used in combination with bleach activators such as tetra-acetylenethylenediamine (TAED), nonanoyloxybenzenesulfonate (NOBS, 5 described in US 4,412,934), 3,5-trimethyl-hexanoloxybenzenesulfonate (ISONOBS, described in EP 120 591) or pentaacetylglucose (PAG), which are perhydrolyzed to form a peracid as the active bleaching species, leading to improved bleaching effect. In addition, very suitable are the bleach 10 activators C8(6-octanamido-caproyl) oxybenzene-sulfonate, C9(6-nonanamido caproyl) oxybenzenesulfonate and C10 (6-decanamido caproyl) oxybenzenesulfonate or mixtures thereof. Also suitable activators are acylated citrate esters such as disclosed in European Patent Application No. 91870207.7.

15 Useful bleaching agents, including peroxyacids and bleaching systems comprising bleach activators and peroxygen bleaching compounds for use in cleaning compositions according to the invention are described in application USSN 08/136,626.

The hydrogen peroxide may also be present by adding an 20 enzymatic system (i.e. an enzyme and a substrate therefore) which is capable of generation of hydrogen peroxide at the beginning or during the washing and/or rinsing process. Such enzymatic systems are disclosed in European Patent Application EP 0 537 381.

25 Bleaching agents other than oxygen bleaching agents are also known in the art and can be utilized herein. One type of non-oxygen bleaching agent of particular interest includes photoactivated bleaching agents such as the sulfonated zinc and/or aluminium phthalocyanines. These materials can be deposited 30 upon the substrate during the washing process. Upon irradiation with light, in the presence of oxygen, such as by hanging clothes out to dry in the daylight, the sulfonated zinc phthalocyanine is activated and, consequently, the substrate is bleached. Preferred zinc phthalocyanine and a photoactivated 35 bleaching process are described in US 4,033,718. Typically, detergent composition will contain about 0.025% to about 1.25%, by weight, of sulfonated zinc phthalocyanine.

Bleaching agents may also comprise a manganese catalyst. The manganese catalyst may, e.g., be one of the compounds

described in "Efficient manganese catalysts for low-temperature bleaching", Nature 369, 1994, pp. 637-639.

Suds suppressors

5 Another optional ingredient is a suds suppressor, exemplified by silicones, and silica-silicone mixtures. Silicones can generally be represented by alkylated polysiloxane materials, while silica is normally used in finely divided forms exemplified by silica aerogels and xerogels and hydrophobic
10 silicas of various types. These materials can be incorporated as particulates, in which the suds suppressor is advantageously releasably incorporated in a water-soluble or waterdispersible, substantially non surface-active detergent impermeable carrier. Alternatively the suds suppressor can be dissolved or dispersed
15 in a liquid carrier and applied by spraying on to one or more of the other components.

A preferred silicone suds controlling agent is disclosed in US 3,933,672. Other particularly useful suds suppressors are the self-emulsifying silicone suds suppressors, described in
20 German Patent Application DTOS 2,646,126. An example of such a compound is DC-544, commercially available from Dow Corning, which is a siloxane-glycol copolymer. Especially preferred suds controlling agent are the suds suppressor system comprising a mixture of silicone oils and 2-alkyl-alkanols. Suitable 2-alkyl-
25 alkanols are 2-butyl-octanol which are commercially available under the trade name Isofol 12 R.

Such suds suppressor system are described in European Patent Application EP 0 593 841.

Especially preferred silicone suds controlling agents are
30 described in European Patent Application No. 92201649.8. Said compositions can comprise a silicone/ silica mixture in combination with fumed nonporous silica such as Aerosil®.

The suds suppressors described above are normally employed at levels of from 0.001% to 2% by weight of the composition,
35 preferably from 0.01% to 1% by weight.

Other components

Other components used in detergent compositions may be employed such as soil-suspending agents, soil-releasing agents,

optical brighteners, abrasives, bactericides, tarnish inhibitors, coloring agents, and/or encapsulated or nonencapsulated perfumes.

Especially suitable encapsulating materials are water soluble capsules which consist of a matrix of polysaccharide and polyhydroxy compounds such as described in GB 1,464,616.

Other suitable water soluble encapsulating materials comprise dextrans derived from ungelatinized starch acid esters of substituted dicarboxylic acids such as described in US 3,455,838. These acid-ester dextrans are, preferably, prepared from such starches as waxy maize, waxy sorghum, sago, tapioca and potato. Suitable examples of said encapsulation materials include N-Lok manufactured by National Starch. The N-Lok encapsulating material consists of a modified maize starch and glucose. The starch is modified by adding monofunctional substituted groups such as octenyl succinic acid anhydride.


Antiredeposition and soil suspension agents suitable herein include cellulose derivatives such as methylcellulose, carboxymethylcellulose and hydroxyethylcellulose, and homo- or co-polymeric polycarboxylic acids or their salts. Polymers of this type include the polyacrylates and maleic anhydride-acrylic acid copolymers previously mentioned as builders, as well as copolymers of maleic anhydride with ethylene, methylvinyl ether or methacrylic acid, the maleic anhydride constituting at least 20 mole percent of the copolymer. These materials are normally used at levels of from 0.5% to 10% by weight, more preferably from 0.75% to 8%, most preferably from 1% to 6% by weight of the composition.

Preferred optical brighteners are anionic in character, examples of which are disodium 4,4'-bis-(2-diethanolamino-4-anilino -s- triazin-6-ylamino)stilbene-2:2' disulphonate, disodium 4, - 4'-bis-(2-morpholino-4-anilino-s-triazin-6-ylamino-stilbene-2:2' - disulphonate, disodium 4,4' - bis-(2,4-dianilino-s-triazin-6-ylamino)stilbene-2:2' - disulphonate, monosodium 4',4'' - bis-(2,4-dianilino-s-tri-azin-6-ylamino)stilbene-2-sulphonate, disodium 4,4' -bis-(2-anilino-4-(N-methyl-N-2-hydroxyethylamino)-s-triazin-6-ylamino)stilbene-2,2' - disulphonate, di-sodium 4,4' -bis-(4-phenyl-2,1,3-triazol-2-yl)-stilbene-2,2' disulphonate, di-so-dium 4,4'bis(2-

anilino-4-(1-methyl-2-hydroxyethylamino)-s-triazin-6-ylamino)stilbene-2,2'-disulphonate, sodium 2(stilbyl-4''-(naphtho-1',2':4,5)-1,2,3, - triazole-2''-sulphonate and 4,4'-bis(2-sulphostyryl)biphenyl.

5 Other useful polymeric materials are the polyethylene glycols, particularly those of molecular weight 1000-10000, more particularly 2000 to 8000 and most preferably about 4000. These are used at levels of from 0.20% to 5% more preferably from 0.25% to 2.5% by weight. These polymers and the previously
10 mentioned homo- or co-polymeric poly-carboxylate salts are valuable for improving whiteness maintenance, fabric ash deposition, and cleaning performance on clay, proteinaceous and oxidizable soils in the presence of transition metal impurities.

Soil release agents useful in compositions of the present
15 invention are conventionally copolymers or terpolymers of terephthalic acid with ethylene glycol and/or propylene glycol units in various arrangements. Examples of such polymers are disclosed in US 4,116,885 and 4,711,730 and EP 0 272 033. A particular preferred polymer in accordance with EP 0 272 033 has
20 the formula:



where PEG is $-(\text{OC}_2\text{H}_4)_n-$, PO is $(\text{OC}_3\text{H}_6\text{O})$ and T is $(\text{pOOC}_6\text{H}_4\text{CO})$.

25 Also very useful are modified polyesters as random copolymers of dimethyl terephthalate, dimethyl sulfoisophthalate, ethylene glycol and 1,2-propanediol, the end groups consisting primarily of sulphobenzoate and secondarily of mono esters of ethylene glycol and/or 1,2-propanediol. The
30 target is to obtain a polymer capped at both end by sulphobenzoate groups, "primarily", in the present context most of said copolymers herein will be endcapped by sulphobenzoate groups. However, some copolymers will be less than fully capped, and therefore their end groups may consist of monoester of
35 ethylene glycol and/or 1,2-propanediol, thereof consist "secondarily" of such species.

The selected polyesters herein contain about 46% by weight of dimethyl terephthalic acid, about 16% by weight of 1,2-propanediol, about 10% by weight ethylene glycol, about 13% by

weight of dimethyl sulfobenzoic acid and about 15% by weight of sulfoisophthalic acid, and have a molecular weight of about 3.000. The polyesters and their method of preparation are described in detail in EP 311 342.

5

Softening agents

Fabric softening agents can also be incorporated into laundry detergent compositions in accordance with the present invention. These agents may be inorganic or organic in type.

10 Inorganic softening agents are exemplified by the smectite clays disclosed in GB-A-1 400898 and in US 5,019,292. Organic fabric softening agents include the water insoluble tertiary amines as disclosed in GB-A1 514 276 and EP 0 011 340 and their combination with mono C_{12} - C_{14} quaternary ammonium salts are
15 disclosed in EP-B-0 026 528 and di-long-chain amides as disclosed in EP 0 242 919. Other useful organic ingredients of fabric softening systems include high molecular weight polyethylene oxide materials as disclosed in EP 0 299 575 and 0 313 146.

20 Levels of smectite clay are normally in the range from 5% to 15%, more preferably from 8% to 12% by weight, with the material being added as a dry mixed component to the remainder of the formulation. Organic fabric softening agents such as the water-insoluble tertiary amines or dilong chain amide materials
25 are incorporated at levels of from 0.5% to 5% by weight, normally from 1% to 3% by weight whilst the high molecular weight polyethylene oxide materials and the water soluble cationic materials are added at levels of from 0.1% to 2%, normally from 0.15% to 1.5% by weight. These materials are
30 normally added to the spray dried portion of the composition, although in some instances it may be more convenient to add them as a dry mixed particulate, or spray them as molten liquid on to other solid components of the composition.

35 Polymeric dye-transfer inhibiting agents

The detergent compositions according to the present invention may also comprise from 0.001% to 10%, preferably from 0.01% to 2%, more preferably from 0.05% to 1% by weight of polymeric dye-transfer inhibiting agents. Said polymeric dye-

transfer inhibiting agents are normally incorporated into detergent compositions in order to inhibit the transfer of dyes from colored fabrics onto fabrics washed therewith. These polymers have the ability of complexing or adsorbing the fugitive dyes washed out of dyed fabrics before the dyes have the opportunity to become attached to other articles in the wash.

Especially suitable polymeric dye-transfer inhibiting agents are polyamine N-oxide polymers, copolymers of N-vinylpyrrolidone and N-vinylimidazole, polyvinylpyrrolidone polymers, polyvinylloxazolidones and polyvinylimidazoles or mixtures thereof.

Addition of such polymers also enhances the performance of the enzymes according to the invention.

The detergent composition according to the invention can be in liquid, paste, gels, bars or granular forms.

Non-dusting granulates may be produced, e.g., as disclosed in US 4,106,991 and 4,661,452 (both to Novo Industri A/S) and may optionally be coated by methods known in the art. Examples of waxy coating materials are poly(ethylene oxide) products (polyethyleneglycol, PEG) with mean molecular weights of 1000 to 20000; ethoxylated nonylphenols having from 16 to 50 ethylene oxide units; ethoxylated fatty alcohols in which the alcohol contains from 12 to 20 carbon atoms and in which there are 15 to 80 ethylene oxide units; fatty alcohols; fatty acids; and mono- and di- and triglycerides of fatty acids. Examples of film-forming coating materials suitable for application by fluid bed techniques are given in GB 1483591.

Granular compositions according to the present invention can also be in "compact form", i.e. they may have a relatively higher density than conventional granular detergents, i.e. from 550 to 950 g/l; in such case, the granular detergent compositions according to the present invention will contain a lower amount of "Inorganic filler salt", compared to conventional granular detergents; typical filler salts are alkaline earth metal salts of sulphates and chlorides, typically sodium sulphate; "Compact" detergent typically comprise not more than 10% filler salt. The liquid compositions according to the present invention can also be in "concentrated form", in such

case, the liquid detergent compositions according to the present invention will contain a lower amount of water, compared to conventional liquid detergents. Typically, the water content of the concentrated liquid detergent is less than 30%, more preferably less than 20%, most preferably less than 10% by weight of the detergent compositions.

The compositions of the invention may for example, be formulated as hand and machine laundry detergent compositions including laundry additive compositions and compositions suitable for use in the pretreatment of stained fabrics, rinse added fabric softener compositions, and compositions for use in general household hard surface cleaning operations and dishwashing operations.

The following examples are meant to exemplify compositions for the present invention, but are not necessarily meant to limit or otherwise define the scope of the invention. In the detergent compositions, the abbreviated component identifications have the following meanings:

- 20 LAS: Sodium linear C_{12} alkyl benzene sulphonate
- TAS: Sodium tallow alkyl sulphate
- XYAS: Sodium $C_{1x} - C_{1y}$ alkyl sulfate
- SS: Secondary soap surfactant of formula 2-butyl octanoic acid
- 25 25EY: A $C_{12} - C_{15}$ predominantly linear primary alcohol condensed with an average of Y moles of ethylene oxide
- 45EY: A $C_{14} - C_{15}$ predominantly linear primary alcohol condensed with an average of Y moles of ethylene oxide
- XYEZS: $C_{1x} - C_{1y}$ sodium alkyl sulfate condensed with an
- 30 average of Z moles of ethylene oxide per mole
- Nonionic: $C_{13} - C_{15}$ mixed ethoxylated/propoxylated fatty alcohol with an average degree of ethoxylation of 3.8 and an average degree of propoxylation of 4.5 sold under the tradename Plurafax LF404 by BASF GmbH
- 35 CFAA: $C_{12} - C_{14}$ alkyl N-methyl glucamide
- TFAA: $C_{16} - C_{18}$ alkyl N-methyl glucamide
- Silicate: Amorphous Sodium Silicate ($SiO_2:Na_2O$ ratio = 2.0)
- NaSKS-6: Crystalline layered silicate of formula $d-Na_2Si_2O_5$
- Carbonate: Anhydrous sodium carbonate

Phosphate: Sodium tripolyphosphate

MA/AA: Copolymer of 1:4 maleic/acrylic acid, average molecular weight about 80,000

Polyacrylate: Polyacrylate homopolymer with an average
5 molecular weight of 8,000 sold under the tradename PA30 by BASF GmbH

Zeolite A: Hydrated Sodium Aluminosilicate of formula $\text{Na}_{12}(\text{AlO}_2\text{SiO}_2)_{12} \cdot 27\text{H}_2\text{O}$ having a primary particle size in the range from 1 to 10 micrometers

10 Citrate: Tri-sodium citrate dihydrate

Citric: Citric Acid

Perborate: Anhydrous sodium perborate monohydrate bleach, empirical formula $\text{NaBO}_2 \cdot \text{H}_2\text{O}_2$

PB4: Anhydrous sodium perborate tetrahydrate

15 Percarbonate: Anhydrous sodium percarbonate bleach of empirical formula $2\text{Na}_2\text{CO}_3 \cdot 3\text{H}_2\text{O}_2$

TAED: Tetraacetyl ethylene diamine

CMC: Sodium carboxymethyl cellulose

DETPMP: Diethylene triamine penta (methylene phosphonic
20 acid), marketed by Monsanto under the Tradename Dequest 2060

PVP: Polyvinylpyrrolidone polymer

EDDS: Ethylenediamine-N, N'-disuccinic acid, [S,S] isomer in the form of the sodium salt

Suds Suppressor: 25% paraffin wax Mpt 50°C, 17% hydrophobic
25 silica, 58% paraffin oil

Granular Suds suppressor: 12% Silicone/silica, 18% stearyl alcohol, 70% starch in granular form

Sulphate: Anhydrous sodium sulphate

HMWPEO: High molecular weight polyethylene oxide

30 TAE 25: Tallow alcohol ethoxylate (25)

Detergent Example I

A granular fabric cleaning composition in accordance with
35 the invention may be prepared as follows:

Sodium linear C_{12} alkyl	6.5
benzene sulfonate	
Sodium sulfate	15.0

	Zeolite A	26.0
	Sodium nitrilotriacetate	5.0
	Enzyme of the invention	0.1
	PVP	0.5
5	TAED	3.0
	Boric acid	4.0
	Perborate	18.0
	Phenol sulphonate	0.1
	Minors	Up to 100

10

Detergent Example II

A compact granular fabric cleaning composition (density 800 g/l) in accord with the invention may be prepared as follows:

15	45AS	8.0
	25E3S	2.0
	25E5	3.0
	25E3	3.0
	TFAA	2.5
20	Zeolite A	17.0
	NaSKS-6	12.0
	Citric acid	3.0
	Carbonate	7.0
	MA/AA	5.0
25	CMC	0.4
	Enzyme of the invention	0.1
	TAED	6.0
	Percarbonate	22.0
	EDDS	0.3
30	Granular suds suppressor	3.5
	water/minors	Up to 100%

Detergent Example III

Granular fabric cleaning compositions in accordance with the invention which are especially useful in the laundering of coloured fabrics were prepared as follows:

	LAS	10.7	-
	TAS	2.4	-
	TFAA	-	4.0

54

	45AS	3.1	10.0
	45E7	4.0	-
	25E3S	-	3.0
	68E11	1.8	-
5	25E5	-	8.0
	Citrate	15.0	7.0
	Carbonate	-	10
	Citric acid	2.5	3.0
	Zeolite A	32.1	25.0
10	Na-SKS-6	-	9.0
	MA/AA	5.0	5.0
	DETPMP	0.2	0.8
	Enzyme of the invention	0.10	0.05
	Silicate	2.5	-
15	Sulphate	5.2	3.0
	PVP	0.5	-
	Poly (4-vinylpyridine)-N-Oxide/copolymer of vinyl-imidazole and vinyl-	-	0.2
20	pyrrolidone		
	Perborate	1.0	-
	Phenol sulfonate	0.2	-
	Water/Minors	Up to 100%	

25 Detergent Example IV

Granular fabric cleaning compositions in accordance with the invention which provide "Softening through the wash" capability may be prepared as follows:

	45AS	-	10.0
30	LAS	7.6	-
	68AS	1.3	-
	45E7	4.0	-
	25E3	-	5.0
	Coco-alkyl-dimethyl hydroxy-	1.4	1.0
35	ethyl ammonium chloride		
	Citrate	5.0	3.0
	Na-SKS-6	-	11.0
	Zeolite A	15.0	15.0
	MA/AA	4.0	4.0

55

	DETPMP	0.4	0.4
	Perborate	15.0	-
	Percarbonate	-	15.0
	TAED	5.0	5.0
5	Smectite clay	10.0	10.0
	HMWPEO	-	0.1
	Enzyme of the invention	0.10	0.05
	Silicate	3.0	5.0
	Carbonate	10.0	10.0
10	Granular suds suppressor	1.0	4.0
	CMC	0.2	0.1
	Water/Minors	Up to 100%	

Detergent Example V

- 15 Heavy duty liquid fabric cleaning compositions in accordance with the invention may be prepared as follows:

		I	II
	LAS acid form	-	25.0
20	Citric acid	5.0	2.0
	25AS acid form	8.0	-
	25AE2S acid form	3.0	-
	25AE7	8.0	-
	CFAA	5	-
25	DETPMP	1.0	1.0
	Fatty acid	8	-
	Oleic acid	-	1.0
	Ethanol	4.0	6.0
	Propanediol	2.0	6.0
30	Enzyme of the invention	0.10	0.05
	Coco-alkyl dimethyl hydroxy ethyl ammonium chloride	-	3.0
	Smectite clay	-	5.0
35	PVP	2.0	-
	Water / Minors	Up to 100%	

THE XYLOGLUCAN SUBSTRATE

In addition to the aforesaid about xyloglucan it should be noted that xyloglucan from tamarind seeds supplied by Megazyme, Ireland has a complex branched structure with glucose, xylose, galactose and arabinose in the ratio of 45:36:16:3. Accordingly, it is strongly believed that an enzyme showing catalytic activity on this xyloglucan also has catalytic activity on other xyloglucan structures from different sources (angiosperms or gymnosperms).

Cotton suspension culture xyloglucan MW 100,000 kDa was obtained from Professor A. Mort of Oklahoma State University. ¹H NMR (D₂O, 80°C) of xyloglucans was used to compare the monosaccharide composition of samples of different origin. The integrals of the anomeric signals from the commercial sample fully agree with the composition given by Megazyme. However, the cotton xyloglucan seems to have a different structure. There appears to be much less galactose and about half of galactose residues are fucosylated. Furthermore, the molar ratio between xylose and glucose is smaller (0.63 compared to 0.77 for the tamarind), which suggest a more open structure of cotton xyloglucan. These findings agree with results obtained with xyloglucan from cotton cells (Buchala et al, Acta Bot. Neerl. 42, 1993, 213-219).

25 Xyloglucan	(Megazyme)	Cotton xyloglucan
Glucose	45 %	52 %
Xylose	35 %	33 %
Galactose	16 %	10 %
Fucose	-	5 %
30 Arabinose	<4 % a	-

a Could not be detected in NMR

MATERIALS AND METHODS

35 Strains

Paenibacillus polymyxa, e.g. *Paenibacillus polymyxa*, ATCC 832, and *Paenibacillus sp.*, DSM 13329, comprises a DNA sequence encoding a xyloglucanase of the invention.

Other strains

E. coli hosts: XL1-Blue MRF⁻ and XL0LR *E. coli* strains were provided by Stratagene Inc. (USA) and used according to the manufacturer's instructions.

5 *B. subtilis* PL1885. (Diderichsen et al., (1990)).

B. subtilis PL1801. This strain is the *B. subtilis* DN1885 where the two major proteases have been inactivated (Diderichsen et al., (1990)).

Competent cells were prepared and transformed as described by
10 Yasbin et al. (1975).

Plasmids

pBK-CAMV: Stratagene inc. La Jolla CA., USA.

Bacteriophage ZAP Express: Stratagene inc. La Jolla CA.,
15 USA.

pMOL944. This plasmid is a pUB110 derivative essentially containing elements making the plasmid propagatable in *Bacillus subtilis*, kanamycin resistance gene and having a strong promoter and signal peptide cloned from the amyL gene of *B. licheniformis*
20 ATCC 14580. The signal peptide contains a SacII site making it convenient to clone the DNA encoding the mature part of a protein in-fusion with the signal peptide. This results in the expression of a Pre-protein which is directed towards the exterior of the cell.

25 The plasmid was constructed by means of ordinary genetic engineering and is briefly described in the following.

Construction of pMOL944:

The pUB110 plasmid (McKenzie, T. et al., 1986,) was digested with the unique restriction enzyme NciI. A PCR fragment
30 amplified from the amyL promoter encoded on the plasmid pDN1981 (Jørgensen et al., 1990) was digested with NciI and inserted in the NciI digested pUB110 to give the plasmid pSJ2624.

The two PCR primers used have the following sequences:

LWN5494 5'-GTCGCCGGGGCGGCCGCTATCAATTGGTAACTGTATCTCAGC -3'
35 # LWN5495 5'-GTCGCCCGGGAGCTCTGATCAGGTACCAAGCTTGTTCGACCTGCAGAA
TGAGGCAGCAAGAAGAT -3'

The primer #LWN5494 inserts a NotI site in the plasmid.

The plasmid pSJ2624 was then digested with SacI and NotI and a new PCR fragment amplified on amyL promoter encoded on the pDN1981 was digested with SacI and NotI and this DNA fragment was inserted in the SacI-NotI digested pSJ2624 to give the
5 plasmid pSJ2670.

This cloning replaces the first amyL promoter cloning with the same promoter but in the opposite direction. The two primers used for PCR amplification have the following sequences:

#LWN5938 5'-GTCGGCGGCCGCTGATCACGTACCAAGCTTGTCTGACCTGCAGAATG
10 AGGCAGCAAGAAGAT -3'
#LWN5939 5'-GTCGGAGCTCTATCAATTGGTAACTGTATCTCAGC -3'

The plasmid pSJ2670 was digested with the restriction enzymes PstI and BclI and a PCR fragment amplified from a cloned DNA sequence encoding the alkaline amylase SP722 (International
15 Patent Application published as WO95/26397 which is hereby incorporated by reference) was digested with PstI and BclI and inserted to give the plasmid pMOL944. The two primers used for PCR amplification have the following sequence:

#LWN7864 5' -AACAGCTGATCACGACTGATCTTTTAGCTTGGCAC-3'
20 #LWN7901 5' -AACTGCAGCCGCGGCACATCATAATGGGACAAATGGG -3'

The primer #LWN7901 inserts a SacII site in the plasmid.

pPL3143 : This plasmid is a pMOL944 derivative in which a terminator has been inserted between the SacII and the NotI site in pMOL944. At the same time a new restriction site for cloning
25 AscI has been inserted.

The plasmid was constructed by means of ordinary genetic engineering and is briefly described in the following.

Construction of pPL3143 : The plasmid pMOL944 was digested with SacII and NotI . A PCR fragment generating a terminator was
30 made using the two primers listed below and plasmid pMOL944 as template. This fragment was digested with EagI and SacII and inserted between the SacII and the NotI site in PMOL944 to create the plasmid pPL3143.

35 Primer 130721 :

5' - CGATCGGCCGATAAAAAACCGGGCGGAAACCGCCCGTCATCTGGCGCGCCTAT
ATACCGCGGCTGCAGAATGAGGCAGCAAG -3'

Primer 130722 :

5' -- GGCGCATTAACGGAATAAAGGGTGT - 3'

Media

TY (as described in Ausubel, F. M. et al. 1995).

LB agar (as described in Ausubel, F. M. et al, 1995).

5 LBPG is LB agar supplemented with 0.5% Glucose and 0.05 M potassium phosphate, pH 7.0

AZCL-xyloglucan is added to LBPG-agar to 0.5 %. AZCL-xyloglucan is from Megazyme, Ireland.

BPX media is described in EP 0 506 780 (WO 91/09129).

10 NZY agar (per liter) 5 g of NaCl, 2 g of MgSO₄, 5 g of yeast extract, 10 g of NZ amine (casein hydrolysate), 15 g of agar; add deionized water to 1 liter, adjust pH with NaOH to pH 7.5 and autoclave

NZY broth (per liter) 5 g of NaCl, 2 g of MgSO₄, 5 g of yeast extract, 10 g of NZ amine (casein hydrolysate); add deionized water to 1 liter, adjust pH with NaOH to pH 7.5 and autoclave

NZY Top Agar (per liter) 5 g of NaCl, 2 g of MgSO₄, 5 g of yeast extract, 10 g of NZ amine (casein hydrolysate), 0.7 % (w/v) agarose; add deionized water to 1 liter, adjust pH with NaOH to pH 7.5 and autoclave.

Xyloglucanase assay (XyloU)

The xyloglucanase activity is measured using AZCL-xyloglucan from Megazyme, Ireland,
25 (<http://www.megazyme.com/purchase/index.html>) as substrate.

A solution of 0.2 % of the blue substrate is suspended in a 0.1 M phosphate buffer pH 7.5 under stirring. The solution is distributed under stirring to 1.5 ml Eppendorf tubes (0.75 ml to each), 50 µl enzyme solution is added and they are incubated in an Eppendorp Thermomixer model 5436 for 20 minutes at 40°C with a mixing of 1200 rpm. After incubation the coloured solution is separated from the solid by 4 minutes centrifugation at 14,000 rpm and the absorbance of the supernatant is measured at 600 nm.

35 One XyloU unit is defined as the amount of enzyme resulting in an absorbance of 0.24 in a 1 cm cuvette at 600 nm.

General molecular biology methods

DNA manipulations and transformations were performed using

standard methods of molecular biology (Sambrook et al. (1989) Molecular cloning: A laboratory manual, Cold Spring Harbor lab., Cold Spring Harbor, NY; Ausubel, F. M. et al. (eds.) "Current protocols in Molecular Biology". John Wiley and Sons, 1995; 5 Harwood, C. R., and Cutting, S. M. (eds.) "Molecular Biological Methods for Bacillus". John Wiley and Sons, 1990).

Enzymes for DNA manipulations were used according to the specifications of the suppliers.

10 The following examples illustrate the invention.

EXAMPLE 1

Cloning of xyloglucanase encoding genes from *Paenibacillus* species

15

Genomic DNA preparation

Strains of *Paenibacillus polymyxa*, including *Paenibacillus polymyxa*, ATCC 842, and the strain *Paenibacillus* sp., DSM 13329, respectively, were propagated in liquid TY medium. After 16 20 hours incubation at 30°C and 300 rpm, the cells were harvested, and genomic DNA isolated by the method described by Pitcher et al. (1989).

Genomic library construction

25 Lambda ZAP libraries were prepared from genomic DNA of the strains *Paenibacillus polymyxa*, *Paenibacillus polymyxa*, ATCC 842, *Paenibacillus* sp., DSM 13329. The ZAP Express cloning kit used was with BamHI digested and dephosphorylated arms from Stratagene. Isolated DNA was partially digested with Sau3A and 30 size fractionated on a 1% agarose gel. DNA was excised from the agarose gel between 3 and 10 Kb and purified using Qiaspin DNA fragment purification procedure (Qiagen GmbH). 100 ng of purified, fractionated DNA was ligated with 1 µg of BamHI dephosphorylated ZAPexpress vector arms (4 degrees overnight). 35 Ligation reaction was packaged directly with GigaPackIII Gold according to the manufacturers instructions (Stratagene). Phage libraries were titered with XL1blue mrf⁻ (Stratagene).

Screening for xyloglucanase clones by functional expression in lambdaZAPExpress

Approximately 10,000 plaque-forming units (pfu) from the genomic library were plated on NZY-agar plates containing 0.1 % AZCL-xyloglucan (MegaZyme, Ireland), using *E. coli* XL1-Blue MRF' (Stratagene, USA) as a host, followed by incubation of the plates at 37°C for 24 hours. Xyloglucanase-positive lambda clones were identified by the formation of blue hydrolysis halos around the positive phage clones. These were recovered from the screening plates by coring the TOP-agar slices containing the plaques of interest into 500 µl of SM buffer and 20 µl of chloroform. The xyloglucanase-positive lambdaZAPExpress clones were plaque-purified by plating an aliquot of the cored phage stock on NZY plates containing 0.1 % AZCL-xyloglucan as above. Single xyloglucanase-positive lambda clones were cored into 500 µl of SM buffer and 20 µl of chloroform, and purified by one more plating round as described above.

Single-clone *in vivo* excision of the phagemids from the xyloglucanase-positive lambdaZAPExpress clones

E. coli XL1-Blue cells (Stratagene, La Jolla Ca.) were prepared and resuspended in 10mM MgSO₄ as recommended by Stratagene (La Jolla, USA). 250µl aliquots of the pure phage stocks from the xyloglucanase-positive clones were combined in Falcon 2059 tubes with 200µl of XL1-Blue MRF' cells (OD₆₀₀=1.0) and >10⁶ pfus/ml of the ExAssist M13 helper phage (Stratagene), and the mixtures were incubated at 37°C for 15 minutes. 3 ml of NZY broth was added to each tube and the tubes were incubated at 37°C for 2.5 hours. The tubes were heated at 65°C for 20 minutes to kill the *E. coli* cells and bacteriophage lambda; the phagemids being resistant to heating. The tubes were spun at 3000 rpm for 15 minutes to remove cellular debris and the supernatants were decanted into clean Falcon 2059 tubes. Aliquots of the supernatants containing the excised single-stranded phagemids were used to infect 200µl of *E. coli* XL0LR cells (Stratagene, OD₆₀₀=1.0 in 10mM MgSO₄) by incubation at 37°C for 15 minutes. 350µl of NZY broth was added to the cells

and the tubes were incubated for 45 min at 37°C. Aliquots of the cells were plated onto LB kanamycin agar plates and incubated for 24 hours at 37°C. Five excised single colonies were re-streaked onto LB kanamycin agar plates containing 0.1 % AZCL-xyloglucan (MegaZyme, Australia). The xyloglucanase-positive phagemid clones were characterized by the formation of blue hydrolysis halos around the positive colonies. These were further analysed by restriction enzyme digests of the isolated plagemid DNA (QiaSpin kit, Qiagen, USA) with EcoRI, PstI, EcoRI-PstI, and HindIII followed by agarose gel electrophoresis.

Nucleotide sequence analysis

The nucleotide sequence of the genomic xyloglucanase clones were determined from both strands by the dideoxy chain-termination method (Sanger et al. (1977)) using 500 ng of Qiaspin-purified template (Qiagen, USA), the Tag deoxy-terminal cycle sequencing kit (Perkin-Elmer, USA), fluorescent labeled terminators and 5 pmol of either pBK-CMV polylinker primers (Stratagene, USA) or custom synthetic oligonucleotide primers. DNA sequence assembly was performed with the DNA Star package (www.DNASTAR.com).

EXAMPLE 2

Subcloning and expression in *B. subtilis* of the xyloglucanase core part of the XYG1006 multidomain gene (hereafter called XYG1006) from *Paenibacillus polymyxa* encoding for the enzyme of the invention

In order to express the xyloglucanase enzyme core part of the XYG1006 multidomain enzyme the following PCR cloning scheme was followed. This cloning makes a translational fusion between the Amylase signal peptide on pPL3143 and the mature part of the XYG1006 multidomain enzyme. At the same time it creates an artificial translational stop codon corresponding to the amino acid number 560 in the XYG1006 multidomain gene.

The XYG1006 encoding DNA sequence was PCR amplified using the PCR primer set consisting of these two oligo nucleotides:

XYG1006 .upper.PstI

5' -GCATTCTGCAGCAGCGGCTGTGGTTCACGGTCAAACGGC -3'

XYG1006 .lower.AscI

5'-GCTAGGCGCGCCTACACTGGAGACGTGTCATTGCCAGTAG -3'

5 Restriction sites PstI and AscI are underlined.

pXYG1006 plasmid DNA from *E.coli*, DSM 13321, was used as template in a PCR reaction using Amplitaq DNA Polymerase (Perkin Elmer) according to manufacturers instructions. The PCR reaction was set up in PCR buffer (10 mM Tris-HCl, pH 8.3, 50 mM KCl, 1.5 mM MgCl₂, 0.01 % (w/v) gelatin) containing 200 µM of each dNTP, 2.5 units of AmpliTaq polymerase (Perkin-Elmer, Cetus, USA) and 100 pmol of each primer.

The PCR reaction was performed using a DNA thermal cycler (Landgraf, Germany). One incubation at 94°C for 1 min followed by thirty cycles of PCR performed using a cycle profile of denaturation at 94°C for 30 sec, annealing at 60°C for 1 min, and extension at 72 °C for 2 min. Five-µl aliquots of the amplification product was analysed by electrophoresis in 0.7 % agarose gels (NuSieve, FMC). The appearance of a DNA fragment size 1,6 kb indicated proper amplification of the gene segment.

Subcloning of PCR fragment:

Fortyfive-µl aliquots of the PCR products generated as described above were purified using QIAquick PCR purification kit (Qiagen, USA) according to the manufacturer's instructions. The purified DNA was eluted in 50 µl of 10mM Tris-HCl, pH 8.5.

5 µg of pPL3143 and twentyfive-µl of the purified PCR fragment was digested with PstI and AscI, electrophoresed in 0.8 % low gelling temperature agarose (SeaPlaque GTG, FMC) gels, the relevant fragments were excised from the gels, and purified using QIAquick Gel extraction Kit (Qiagen, USA) according to the manufacturer's instructions. The isolated PCR DNA fragment was then ligated to the PstI-AscI digested and purified pPL3143. The ligation was performed overnight at 16°C using 0.5 µg of each DNA fragment, 1 U of T4 DNA ligase and T4 ligase buffer (Boehringer Mannheim, Germany).

The ligation mixture was used to transform competent *B.subtilis* PL1801 cells. The transformed cells were plated onto

LBPG agar plates containing 10 µg/ml of Kanamycin and 0,2% AZCL-Xyloglucan (Megazyme). After 18 hours incubation at 37°C colonies were seen on plates. Several clones showing a blue halo around the colony were analyzed by isolating plasmid DNA from overnight culture broth.

One such positive clone was re-streaked several times on agar plates as used above, this clone was called PL3344. The clone PL3344 was grown overnight in TY-10 µg/ml Kanamycin at 37°C. The following day, 1 ml of cells were used to isolate plasmid from the cells using the Qiaprep Spin Plasmid Miniprep Kit #27106 according to the manufacturers recommendations for *B. subtilis* plasmid preparations. This DNA was DNA sequenced and revealed the DNA sequence corresponding to the mature part of the *Paenibacillus polymyxa* xyloglucanase encoding domain correctly inserted into the pPL3143 cloning vector (Domain 1-1680 of SEQ ID NO:1. The *Bacillus subtilis* strain PL3344 is thus containing a plasmid enabling the expression of the xyloglucanase domain of the XYG1006 multi domain gene. The translation product is thus from amino acid number 1 to 559 (SEQ ID no:2) which after cleavage of the signal peptide is expected to be secreted from *B. subtilis* as a protein spanning from amino acid 36 to 559 (SEQ ID no:2).

SOMEBODY PLEASE EXPLAIN WHY KkSc POLYPEPTIDE STARTS AT POSITION 35, PLJ'S POLYPEPTIDE STARTS AT POSITION 36, MaS ENZYME STARTS AT POSITION 40 (I HAVE CHOSEN POSITION 40 AS STARTING AA, SINCE THE XYLOGLUCANASE 40-559 DOES PERFORM)? I cannot answer this, ask Per and Martin

PL3344 was grown in 25 x 200 ml BPX media with 10 µg/ml of Kanamycin in two 500 ml baffled shake flasks for 5 days at 37°C at 300 rpm.

EXAMPLE 3

Purification and characterization of xyloglucanase from *Paenibacillus polymyxa*

The clone XYG1006 obtained as described in example 2 (PL3344 expressed in *B. subtilis*) was incubated in 4500 ml of PS-1 containing mg/ml kanamycin from shake flasks with a final pH of 5.5.

The fermentation medium was flocculated using cationic

flocculation agent C521 (10% solution) and 0.1% solution of anionic agent A130: To 4500 ml of broth, 200 ml of C521 was added (10%) simultaneously with 400 ml of A130 (0.1%) under constant stirring at room temperature. The flocculated material
5 was separated by centrifugation using a Sorval RC 3B centrifuge at 4,500 rpm for 30 minutes. The supernatant was adjusted to pH 7.0 using sodium hydroxide and then batch treated with 300 g of HPQ Sepharose equilibrated with 50 mM tris pH 7.0, and the unbound material was filter sterilized through a 0.7 μ m filter.

10 The liquid was concentrated into 550 ml using filtron ultrafiltration with a MW cut off of 10 kDa. The concentrate was then diluted to 900 ml and passed over a S-Sepharose column equilibrated with 25 mM sodium acetate pH 5.0, and the not bound material was concentrated to 390 ml containing 370,000 XyloU.

15 REMOVE THIS TOO!

For obtaining a pure enzyme 2 ml of this partial pure enzyme was applied to a size chromatography (Superdex 75) column equilibrated with 0.1 M Sodium acetate pH 6.0. The xyloglucanase eluted as a single peak with a MW of 58 kDa in SDS-PAGE and with
20 a specific activity of 255 XyloU units per mg protein.

The cloned xyloglucanase of the invention was used for raising rabbit antiserum.

After electroblotting of this band the N-terminal was determined as:

25 TAKTITIKVDTFKDRK

This is in agreement with the amino acid sequence shown in SEQ ID NO:2 deduced from the DNA sequence shown in SEQ ID NO:1 with a 40 amino acid pro sequence. The calculated MW from the deduced sequence was 58 kDa and the calculated pI was 5.7. The
30 molar extinction coefficient at 280 nm was 105,640.

The enzyme melted in the DSC (Differential Scanning Calorimeter) at 61.8°C at pH 6.0.

The xyloglucanase showed optimal activity at 50° at pH 7.5 using the xylounits (XyloU) assay at different temperatures.

35 The xyloglucanase was more than 30% active between pH 5.0 and 8.0.

Kinetic determinations were performed using soluble tamarind gum xyloglucan at pH 7.5 and at 40°C, 20 minutes incubation time and measurement of the formation of reducing

ends. The kCat of 220 per sec was determined. The kM was 0.2 g/l. The Michaelis-Menten kinetic values could be determined.

EXAMPLE 4

5 Stability and activity of *Paenibacillus polymyxa* xyloglucanase in commercial liquid detergents

The xyloglucanase characterized in Example 3 was fully stable between pH 5 and 10 at room temperature, and had a half life of more than 50 days when incubated in a full formulated US liquid detergent as US Tide at 30° C. The xyloglucanase was fully active in US liquid Tide and in European liquid Ariel, using 20 min. incubation at 40°C.

LITERATURE

Ausubel, F. M. et al. (eds.) "Current protocols in Molecular Biology". John Wiley and Sons, 1995.

5

N. C. Carpita and D. M. Gibeaut (1993) *The Plant Journal* 3:1-30.

T. Christensen et al. *Biotechnology* vol 6 page 1419-1422, 1988.

10 Devereux et al. (1984) *Nucleic Acids Res.* 12, 387-395.

Diderichsen, B., Wedsted, U., Hedegaard, L., Jensen, B. R., Sjøholm, C. (1990) Cloning of aldB, which encodes alpha-acetolactate decarboxylase, an exoenzyme from *Bacillus brevis*.

15 *J. Bacteriol.* 172:4315-4321.

Dretzen, G., Bellard, M., Sassone-Corsi, P., Chambon, P. (1981) A reliable method for the recovery of DNA fragments from agarose and acrylamide gels. *Anal. Biochem.*, 112, 295-298.

20

Eriksson, O.E. & Hawksworth, D.L.: *Systema Ascomycetum* vol 12 (1993).

S. C. Fry et al (1992) *Biochemical Journal* 282:821-828

25

Hawksworth, D.L., Kirk, P.M., Sutton, B.C. and Pegler, D.N.: *Dictionary of the fungi*, International Mycological Institute, 616 pp (1995);

30 T. Hayashi and D. P. Delmer (1988) *Carbohydrate Research* 181:273-277.

Henrissat, B. 1991. A classification of glycosyl hydrolases based on amino acid sequence similarities. *Biochem. J.*,

35 280:309-316.

Henrissat, B., and A. Bairoch. 1993. New families in the classification of glycosyl hydrolases based on amino acid sequence similarities. *Biochem. J.*, 293:781-788.

Jülich, W.: Higher Taxa of Basidiomycetes, *Bibliotheca Mycologia* 85, 485 pp (1981).

5 Jørgensen, P.L. et al., 1990, *Gene*, 96, p. 37-41.

McKenzie, T. et al., 1986, *Plasmid* 15:93-103.

Leatherbarrow, R. J. (1992) *Grafit* version 3.0 Erithacus
10 Software Ltd. Staines, U.K.

Lever, M. (1972) A new reaction for colormetric determination of carbohydrates. *Anal. Biochem.* 47, 273-279.

15 O'Donnell, K.: *Zygomycetes in culture*, University of Georgia, US, 257 pp (1979).

Pitcher, D. G., Saunders, N. A., Owen, R. J. (1989). Rapid extraction of bacterial genomic DNA with guanidium thiocyanate.
20 *Lett. Appl. Microbiol.*, 8, 151-156.

J. K. C. Rose et al (1996) *Plant Physiology* 110:493-499

Sanger, F., Nicklen, S., and Coulson, A. R. (1977) *Proc. Natl.*
25 *Acad. Sci. U. S. A.* 74, 5463-5467

A.L. Sonenshein, J.A. Hoch and Richard Losick (Eds.) (1993) *Bacillus subtilis* and other Gram-Positive Bacteria, American Society for microbiology, p.618.

30

Vincken, J.P., Beldman, G., and Voragen, A.G.J. Substrate-specificity of endoglucanases - what determines xyloglucanase activity. *Carbohydrate Research* 298(4):299-310, 1997.

35 Von Arx, J. A.: The genera of fungi sporulating in culture, 424 pp (1981).

R. L. Whistler and J. N. BeMiller (1993) *Industrial gums: Polysaccharides and their derivatives*, Academic Press Inc.

Yasbin, R.E., Wilson, G.A. and Young, F.E. (1975) Transformation and transfection in lysogenic strains of *Bacillus subtilis* : evidence for selective induction of prophage in competent cells.

5 J. Bacteriol, 121:296-304.

W.S. York et al (1996) Carbohydrate Research 285:99-128.

SEQUENCE LISTING

<110> NOVO NORDISK A/S

5 <120> FAMILY 44 XYLOGLUCANASES

<130> 10017

<140>

10 <141>

<160> 6

<170> PatentIn Ver. 2.1

15

<210> 1

<211> 4059

<212> DNA

<213> Paenibacillus polymyxa

20

<400> 1

	atgagggcga	aaaatagtag	taatcttttg	ttcaaacgtt	ccaaatggct	gcctgtcgtc	60
	atggcctgca	cgatgatagt	agggggggct	ttacctgctc	cagctgtggt	tcacgggtcaa	120
	acggcaaaga	ctattactat	taaagtagat	acattcaagg	atcgtaagcc	tatttagccct	180
25	tatatatacg	gtacaaatca	ggatttgcca	ggcgtatgaa	atatggctgc	cagacgactt	240
	ggtggcaacc	gaatgaccgg	atacaactgg	gaaaacaata	tgtccaatgc	aggaagtgc	300
	tggcagcaat	ctagcgataa	ctatttatgc	agtaatgggt	gcctgacaca	agccgaatgt	360
	gaaaagccag	gagcgggtgac	gacttcgttt	catgaccaat	cgctgaagct	tggcacttat	420
	tctttagtta	cggtgccgat	ggccgggttat	gtggctaagg	atggaaacgg	aagtgtgcag	480
30	gaaagcgaaa	aggcccccttc	cgctcggttg	aatcaggctc	taaaacgcaa	aaatgcaccg	540
	ttccaactac	agcctgatct	gaatgacaat	cgggtctatg	tggatgagtt	cgctcatttt	600
	ttagtgaaca	agtacggcac	tgcttcaaca	aaggcggggg	tgaaaggata	tgccctcgac	660
	aatgaacccg	ctctctggtc	gcatacgcac	ccacgcattc	atggtgaaaa	agtcggagcg	720
	aaagagttgg	tagaccggtc	agtcagttta	tccaaagctg	tgaaagcgat	tgacgcgggg	780
35	gcagaggttt	ttggcccggg	tctttacgga	ttggcgccct	ataaagatct	tcaaactgca	840
	cctgattgctc	actctgttaa	aggcaattat	agctgggttc	tagactatta	cctggatcaa	900
	atgcgcttta	gctcgcaagt	cgaaggcaag	agattgctgg	atgtattcga	cgtaactctg	960
	tatcccgaag	cgatggcgcg	aggcatacga	attacgaatg	aggtaggcaa	tgacgaaacg	1020
	aagaaagcca	gaatgcaggc	acctcgccac	ttgtgggacc	cgacctataa	ggaagatagt	1080
40	tggatcgctc	aatggaaacag	cgagtttttg	cccatactac	ctcgattgaa	gcagtcgggtg	1140
	gataaatatt	atccgggaac	caagctggca	atgaccgagt	atagctatgg	cgccgaaaaat	1200
	gatattttccg	gcgggattgc	gatgaccgat	gtgctgggta	tcttggggcaa	aaatgatggt	1260
	tatatggcaa	actactggaa	gctaaaggat	gggtgtcaaca	actacgttag	tgccgcttac	1320
	aagctttatc	gcaattatga	cggaaaaaac	tctactttcg	gtgataccag	tgtagtgctg	1380
45	caaacatcgg	atattgtcaa	tagctcggtc	catgcttctg	taacgaatgc	atccgacaaa	1440
	gaactgcctc	tcgttgtcat	gaataaaaagc	atggacagcg	cattcgacgc	ccaatttgat	1500
	ctttccggcg	cgaagactta	catttcgggt	aaagtatggg	ggttcgataa	aaacagctcg	1560
	caaataaag	aagcagcgcc	aatcacgcaa	atttcaggca	accgttttac	ttataccgta	1620
	ccgcctttga	cggcatatca	cattgtgctg	actactggca	atgacacgtc	tccagtgga	1680
50	ggtcctgaaa	gctttaagct	gaaagctgag	gctgggtgat	ggaaagtcca	tttatcctgg	1740
	gatgcttcca	gcggagttgt	aggatacagc	gtacagcggg	caacagatga	aaacggccct	1800
	ttcactgctg	tagcatccaa	cttgaccgaa	acgtcttata	cggatactaa	cgtagacaa	1860
	ggtacttcat	actattacaa	agtaaccgcc	aaaaccaata	agggatcgag	cgaatccaat	1920
	attttgaaa	cggttcgcaa	gatgcctgta	aacgggtccc	ctcgctatga	agccgaagaa	1980
55	ggcacgctga	agggaaacat	tgtggaatcc	agcgggaccg	gctactccgg	tgctgggtat	2040
	gtaacgaatt	tccacaatcc	aggggattct	ctgacgatga	cgattcaggc	tcccacggca	2100
	ggcttgta	atcttacaat	cggctaccgt	tctcctcatg	atgacaaaac	caccaatttc	2160
	tcattaaacg	gcaaagcggt	tggcgaactg	ctgcttaaga	aaacggctga	ttttaaagaa	2220
	acttcgggag	gcaaggtgct	gttgaatgca	ggcggaata	cgatcagttt	tgaaacaggc	2280
60	tggggctggg	acgatatcga	ctacgtcaga	ctggagcctg	ccgctgaccg	cccacctcat	2340
	gcggtaacca	aaacgcttac	caatccgaat	gcgacggtag	aagcaaaagc	attgatgaac	2400
	tatctgggtg	atcaatacgg	gaagaatatg	ctctctgggt	aagaggaaat	aaacgaat	2460
	gattggcttc	aagccaatgt	aggtaaaaag	ccggcgattg	cagcgcttga	cctgatcgac	2520
	tattcgccaa	gcagagcgga	acacgggtct	agttccacag	aggcagaaaa	ggcgattgca	2580
65	tgggataaag	aaggggggat	cgttaccttt	gcatggcact	ggaacgcacc	gaaaggtctg	2640
	atcgatacgc	agggaaaaga	atgggtggaga	ggcttctatg	ccgattcaac	cacattcgat	2700
	atagaatatg	cgatgaatca	tccagagtcc	gaagattata	aattacttat	tcgcgacatc	2760
	gatgtgattg	cagggcaatt	gaagaagttg	caggatgcga	aggttccctgt	cctgttccgt	2820
	cctttgcacg	aagcggaagg	aaaatggttc	tggtggggcg	ccaaagggtcc	tgagcctgtt	2880
70	aaaaagctgt	atattttaat	gcacgaccgt	ttgacgaatg	tgcacaaatt	gaacaatctg	2940

```

atttgggtat ggaattctgt tgctccggat tggatccgg gagacgagta tgtggatatt 3000
ttgagctttg actcttatcc gcaagcaggt gattacagcc cgcaaatttc aaaatacgaa 3060
gaccttggtg cattgggcaa ggacaaaaag ctagttgcca tgagcgaaaa tggaccgatc 3120
ccggaccctg atttgatgaa ggcgtatcaa gctcattgga gctgggtcgc tacatgggat 3180
5 ggagattttg tgagagacgg caaacaaaac agccttgagc atctgaaaaa agtgtataat 3240
catccgaacg tcattacgct ggatgagctc ccaacgaact taaaaacgta tggcattact 3300
gagcagccgt ccgtaccggg cagcttcacg ctgaacgctg cgggtgaaac ggcgaaagta 3360
tcgctaagct ggacagcatc ggcgaatgca aaaagctatg aagtgaagcg ttcgacgact 3420
gaaaacggcg cgttcgccac tgtagcgagt gatgtatatg gcagtagcta caccgacaca 3480
10 gctgtaacgg cagatacgac gtactactac caagtcgtag cgaagaacga tgcaggacag 3540
acgctgtcga acacgggctag cgcaatgccg aaagcggata ctcagcagcc gacgacagga 3600
ctgctgctcc aatatcgcac agcagatact aaggtgaacg ataatacact caatccgcaa 3660
ttccaaattg taaacaaagg cacaacctcc ataccgatca acgagttgaa aattcgctac 3720
tactacacaa tcgacgggtga ccgtgagcag actttcaact gcgactatgc gacgctgagc 3780
15 tgctcaaagc tgaacggtaa actgggttaa atggagaagg ctgcaaccgg tgccgattat 3840
tatttggaag ccagtttcaa ttcggatgca ggcgtgttag cacctggagg aagcacgggc 3900
gatatccaaa cccgtattca taagacagac tggtcgaact ataacgaaag tgacgattat 3960
tcgtataaag gcacgcaaac ctcatttgcc gatcatccta aagttacctt gtatcataac 4020
ggcgtacttg tttggggaac cgagccgaca gctaattaa 4059

```

```

20 <210> 2
    <211> 1352
    <212> PRT
25 <213> Paenibacillus polymyxa

```

```

<400> 2
Met Arg Ala Lys Asn Ser Ser Asn Leu Leu Phe Lys Arg Ser Lys Trp
1      5      10      15
30 Leu Pro Val Val Met Ala Cys Thr Met Ile Val Gly Gly Ala Leu Pro
    20      25      30
35 Ala Pro Ala Val Val His Gly Gln Thr Ala Lys Thr Ile Thr Ile Lys
    35      40      45
Val Asp Thr Phe Lys Asp Arg Lys Pro Ile Ser Pro Tyr Ile Tyr Gly
    50      55      60
40 Thr Asn Gln Asp Leu Ala Gly Asp Glu Asn Met Ala Ala Arg Arg Leu
    65      70      75      80
Gly Gly Asn Arg Met Thr Gly Tyr Asn Trp Glu Asn Asn Met Ser Asn
    85      90      95
45 Ala Gly Ser Asp Trp Gln Gln Ser Ser Asp Asn Tyr Leu Cys Ser Asn
    100      105      110
Gly Gly Leu Thr Gln Ala Glu Cys Glu Lys Pro Gly Ala Val Thr Thr
50      115      120      125
Ser Phe His Asp Gln Ser Leu Lys Leu Gly Thr Tyr Ser Leu Val Thr
    130      135      140
55 Leu Pro Met Ala Gly Tyr Val Ala Lys Asp Gly Asn Gly Ser Val Gln
    145      150      155      160
Glu Ser Glu Lys Ala Pro Ser Ala Arg Trp Asn Gln Val Val Asn Ala
    165      170      175
60 Lys Asn Ala Pro Phe Gln Leu Gln Pro Asp Leu Asn Asp Asn Arg Val
    180      185      190
Tyr Val Asp Glu Phe Val His Phe Leu Val Asn Lys Tyr Gly Thr Ala
65      195      200      205
Ser Thr Lys Ala Gly Val Lys Gly Tyr Ala Leu Asp Asn Glu Pro Ala
    210      215      220
70 Leu Trp Ser His Thr His Pro Arg Ile His Gly Glu Lys Val Gly Ala

```


	225	230	235	240
	Lys Glu Leu Val Asp Arg Ser Val Ser Leu Ser Lys Ala Val Lys Ala			
	245		250	255
5	Ile Asp Ala Gly Ala Glu Val Phe Gly Pro Val Leu Tyr Gly Phe Gly			
	260	265	270	
10	Ala Tyr Lys Asp Leu Gln Thr Ala Pro Asp Trp Asp Ser Val Lys Gly			
	275	280	285	
	Asn Tyr Ser Trp Phe Val Asp Tyr Tyr Leu Asp Gln Met Arg Leu Ser			
	290	295	300	
15	Ser Gln Val Glu Gly Lys Arg Leu Leu Asp Val Phe Asp Val His Trp			
	305	310	315	320
	Tyr Pro Glu Ala Met Gly Gly Gly Ile Arg Ile Thr Asn Glu Val Gly			
	325	330	335	
20	Asn Asp Glu Thr Lys Lys Ala Arg Met Gln Ala Pro Arg Thr Leu Trp			
	340	345	350	
	Asp Pro Thr Tyr Lys Glu Asp Ser Trp Ile Ala Gln Trp Asn Ser Glu			
25	355	360	365	
	Phe Leu Pro Ile Leu Pro Arg Leu Lys Gln Ser Val Asp Lys Tyr Tyr			
	370	375	380	
30	Pro Gly Thr Lys Leu Ala Met Thr Glu Tyr Ser Tyr Gly Gly Glu Asn			
	385	390	395	400
	Asp Ile Ser Gly Gly Ile Ala Met Thr Asp Val Leu Gly Ile Leu Gly			
	405	410	415	
35	Lys Asn Asp Val Tyr Met Ala Asn Tyr Trp Lys Leu Lys Asp Gly Val			
	420	425	430	
	Asn Asn Tyr Val Ser Ala Ala Tyr Lys Leu Tyr Arg Asn Tyr Asp Gly			
40	435	440	445	
	Lys Asn Ser Thr Phe Gly Asp Thr Ser Val Ser Ala Gln Thr Ser Asp			
	450	455	460	
45	Ile Val Asn Ser Ser Val His Ala Ser Val Thr Asn Ala Ser Asp Lys			
	465	470	475	480
	Glu Leu His Leu Val Val Met Asn Lys Ser Met Asp Ser Ala Phe Asp			
	485	490	495	
50	Ala Gln Phe Asp Leu Ser Gly Ala Lys Thr Tyr Ile Ser Gly Lys Val			
	500	505	510	
	Trp Gly Phe Asp Lys Asn Ser Ser Gln Ile Lys Glu Ala Ala Pro Ile			
55	515	520	525	
	Thr Gln Ile Ser Gly Asn Arg Phe Thr Tyr Thr Val Pro Pro Leu Thr			
	530	535	540	
60	Ala Tyr His Ile Val Leu Thr Thr Gly Asn Asp Thr Ser Pro Val Glu			
	545	550	555	560
	Gly Pro Glu Ser Phe Lys Leu Lys Ala Glu Ala Gly Asp Gly Lys Val			
	565	570	575	
65	His Leu Ser Trp Asp Ala Ser Ser Gly Val Val Gly Tyr Ser Val Gln			
	580	585	590	
	Arg Ala Thr Asp Glu Asn Gly Pro Phe Thr Ala Val Ala Ser Asn Leu			
70	595	600	605	

Thr Glu Thr Ser Tyr Thr Asp Thr Asn Val Thr Asn Gly Thr Ser Tyr
 610 615 620
 5 Tyr Tyr Lys Val Thr Ala Lys Thr Asn Lys Gly Ser Ser Glu Ser Asn
 625 630 635 640
 Ile Leu Lys Ala Val Pro Lys Met Pro Val Asn Gly Pro Ala Arg Tyr
 645 650 655
 10 Glu Ala Glu Glu Gly Thr Leu Lys Gly Thr Ile Val Glu Ser Ser Gly
 660 665 670
 Thr Gly Tyr Ser Gly Ala Gly Tyr Val Thr Asn Phe His Asn Pro Gly
 675 680 685
 15 Asp Ser Leu Thr Met Thr Ile Gln Ala Pro Thr Ala Gly Leu Tyr Asn
 690 695 700
 20 Leu Thr Ile Gly Tyr Arg Ser Pro His Asp Asp Lys Arg Thr Asn Phe
 705 710 715 720
 Ser Leu Asn Gly Lys Ala Phe Gly Glu Leu Leu Leu Lys Lys Thr Ala
 725 730 735
 25 Asp Phe Lys Glu Thr Ser Gly Gly Lys Val Leu Leu Asn Ala Gly Ala
 740 745 750
 Asn Thr Ile Ser Phe Glu Thr Gly Trp Gly Trp Tyr Asp Ile Asp Tyr
 755 760 765
 30 Val Arg Leu Glu Pro Ala Ala Asp Arg Pro Pro His Ala Val Thr Lys
 770 775 780
 35 Thr Leu Thr Asn Pro Asn Ala Thr Val Glu Ala Lys Ala Leu Met Asn
 785 790 795 800
 Tyr Leu Val Asp Gln Tyr Gly Lys Asn Met Leu Ser Gly Gln Glu Glu
 805 810 815
 40 Ile Asn Glu Ile Asp Trp Leu Gln Ala Asn Val Gly Lys Lys Pro Ala
 820 825 830
 Ile Ala Ala Leu Asp Leu Ile Asp Tyr Ser Pro Ser Arg Ala Glu His
 835 840 845
 45 Gly Leu Ser Ser Thr Glu Ala Glu Lys Ala Ile Ala Trp Asp Lys Gln
 850 855 860
 50 Gly Gly Ile Val Thr Phe Ala Trp His Trp Asn Ala Pro Lys Gly Leu
 865 870 875 880
 Ile Asp Thr Gln Gly Lys Glu Trp Trp Arg Gly Phe Tyr Ala Asp Ser
 885 890 895
 55 Thr Thr Phe Asp Ile Glu Tyr Ala Met Asn His Pro Glu Ser Glu Asp
 900 905 910
 Tyr Lys Leu Leu Ile Arg Asp Ile Asp Val Ile Ala Gly Gln Leu Lys
 915 920 925
 60 Lys Leu Gln Asp Ala Lys Val Pro Val Leu Phe Arg Pro Leu His Glu
 930 935 940
 65 Ala Glu Gly Lys Trp Phe Trp Trp Gly Ala Lys Gly Pro Glu Pro Val
 945 950 955 960
 Lys Lys Leu Tyr Ile Leu Met His Asp Arg Leu Thr Asn Val His Lys
 965 970 975
 70

Leu Asn Asn Leu Ile Trp Val Trp Asn Ser Val Ala Pro Asp Trp Tyr
 980 985 990
 5 Pro Gly Asp Glu Tyr Val Asp Ile Leu Ser Phe Asp Ser Tyr Pro Gln
 995 1000 1005
 Ala Gly Asp Tyr Ser Pro Gln Ile Ser Lys Tyr Glu Asp Leu Val Ala
 1010 1015 1020
 10 Leu Gly Lys Asp Lys Lys Leu Val Ala Met Ser Glu Asn Gly Pro Ile
 1025 1030 1035 1040
 Pro Asp Pro Asp Leu Met Lys Ala Tyr Gln Ala His Trp Ser Trp Phe
 1045 1050 1055
 15 Ala Thr Trp Tyr Gly Asp Phe Val Arg Asp Gly Lys Gln Asn Ser Leu
 1060 1065 1070
 Glu His Leu Lys Lys Val Tyr Asn His Pro Asn Val Ile Thr Leu Asp
 1075 1080 1085
 Glu Leu Pro Thr Asn Leu Lys Thr Tyr Gly Ile Thr Glu Gln Pro Ser
 1090 1095 1100
 25 Val Pro Gly Ser Phe Thr Leu Asn Ala Ala Gly Glu Thr Ala Lys Val
 1105 1110 1115 1120
 Ser Leu Ser Trp Thr Ala Ser Ala Asn Ala Lys Ser Tyr Glu Val Lys
 1125 1130 1135
 30 Arg Ser Thr Thr Glu Asn Gly Ala Phe Ala Thr Val Ala Ser Asp Val
 1140 1145 1150
 Tyr Gly Ser Ser Tyr Thr Asp Thr Ala Val Thr Ala Asp Thr Thr Tyr
 1155 1160 1165
 Tyr Tyr Gln Val Val Ala Lys Asn Asp Ala Gly Gln Thr Leu Ser Asn
 1170 1175 1180
 40 Thr Ala Ser Ala Met Pro Lys Ala Asp Thr Gln Gln Pro Thr Thr Gly
 1185 1190 1195 1200
 Leu Leu Leu Gln Tyr Arg Thr Ala Asp Thr Lys Val Asn Asp Asn His
 1205 1210 1215
 45 Leu Asn Pro Gln Phe Gln Ile Val Asn Lys Gly Thr Thr Ser Ile Pro
 1220 1225 1230
 Ile Asn Glu Leu Lys Ile Arg Tyr Tyr Tyr Thr Ile Asp Gly Asp Arg
 1235 1240 1245
 Glu Gln Thr Phe Asn Cys Asp Tyr Ala Thr Leu Ser Cys Ser Lys Leu
 1250 1255 1260
 55 Asn Gly Lys Leu Val Lys Met Glu Lys Ala Ala Thr Gly Ala Asp Tyr
 1265 1270 1275 1280
 Tyr Leu Glu Val Ser Phe Asn Ser Asp Ala Gly Val Leu Ala Pro Gly
 1285 1290 1295
 60 Gly Ser Thr Gly Asp Ile Gln Thr Arg Ile His Lys Thr Asp Trp Ser
 1300 1305 1310
 Asn Tyr Asn Glu Ser Asp Asp Tyr Ser Tyr Lys Gly Thr Gln Thr Ser
 1315 1320 1325
 Phe Ala Asp His Pro Lys Val Thr Leu Tyr His Asn Gly Val Leu Val
 1330 1335 1340
 70 Trp Gly Thr Glu Pro Thr Ala Asn

1345

1350

<210> 3

5 <211> 4056

<212> DNA

<213> *Paenibacillus polymyxa*

<400> 3

10	atgaaggcga	aaaatagtag	tagtatttgg	tccaaacggt	ccaaatggct	gcctgtcgtc	60
	atggcatgca	cgattatagt	agggggtgct	ctaccgactc	caactgtagt	tcacgggtcaa	120
	acggcaaaga	ctgttaccat	taaagtcgat	acatccaagg	atcgtaagcc	tatttagccct	180
	tatatattacg	gtacgaatca	ggagttggca	ggcgatgaga	atctgactgc	cagacgactt	240
	ggtggcaatc	gaatgaccgg	atataactgg	gaaaacaata	tgtccaatgc	aggaagcgac	300
15	tggatgcagt	ccagcgatag	ctatttatgc	gacaacgccc	gattgacaaa	agccgaatgt	360
	gaaaagccag	gtgcggtggc	aacctcgttt	cacgatcaat	cgctgaagca	gggcacatat	420
	tcttttagtca	cactgccgat	ggccggttat	gtggccaagg	atggaaacgg	aagtgtgcag	480
	gaaagcgaaa	aggctccttc	cgctcggtgg	aatgaggtcg	taaacgctaa	aaatgcgccc	540
	tttcaattgc	agcctgatct	gaaagacaat	caggtttatg	cggatgaatt	cgtcaacttt	600
20	ttagtgaaaa	agtacggcgt	tgcttcaaca	aaaacggggc	tgaaaggata	ctcgctcgac	660
	aatgaacccg	ctctctggtc	gcatacgcat	ccgcgcattc	atggtgaaaa	ggtcggagcg	720
	aaagagttgg	tagaccggtc	ggtaagttta	tccaaagccg	ctaaggcggt	tgacgcgggt	780
	cgggaaattt	ttgggcccgt	tctttacggt	tttggcgcc	ataaagatct	tcaaactgca	840
	octgattgga	actctgtaaa	aggcaactac	agctgggtcg	tggactatta	cctcgatcaa	900
25	atgcgcctca	gctcgcaagc	cgaaggcaag	agattgctgg	atgtcttcga	tgtacactgg	960
	tatcctgaag	cgatggggcg	aggcatacga	attacaaatg	aggtaggcaa	cgacgaaacg	1020
	aagaaagcca	gaatgcaagc	gcctcgctac	ttgtgggatc	cgacctacaa	ggaagatagc	1080
	tggatcgctc	aatggaacag	tgaattcttg	cctttactgc	ctcgattaaa	gcagtcggtg	1140
	gataagtatt	acccgggaac	caagctggct	ttgactgagt	atagctatgg	tggcgaaaat	1200
30	gatatttccg	gcggtatcgc	tatggccgat	gtgctgggca	tcttgggcaa	aaacgacgtt	1260
	tatatggcaa	actactggaa	gttaaaagat	ggtgccaaca	actacgttag	tgccgcttac	1320
	aagctttacc	cgcaattatga	cggaaaaagc	ctctacttcg	gtgatatcag	cgttcatgcg	1380
	caaacgtcgg	atattgttaa	tagctcgggt	catgcttcgg	taacggatgc	atcctacaaa	1440
	gaactgcacc	tcggtgtcat	gaataaaaagc	atggacagtg	cattcgacgc	ccaatttgat	1500
35	ctttccggcg	agacgactta	cggttccggg	aaagtattgg	gtttcgacaa	aatagctcgt	1560
	caaattaagg	aagcagcgcc	aatcacscaa	atctcaggca	accgytttac	ctatacagta	1620
	ccgcctttga	cggcttatca	catcgtgttg	actgccggca	atgatacacc	tgtagaaaat	1680
	cctgaaagct	ttgcgctgag	ggctgaggct	ggcgatggaa	agtcgattta	tctggacgct	1740
	tccagcggag	ttgtaggtta	cagcgtacag	cgggcaacgt	atgaaaacgg	tccttttgct	1800
40	gctgtagcat	ccaacttggg	cgaacgctct	tatacggata	cgaacgtaac	gaacggcact	1860
	tcttattatt	ataaaaataac	cgcaaaaaaca	aagacgggaa	cgagcgcac	caatgtcttg	1920
	aaagcgggtc	cgcggggcgcc	tgtagacggg	ccggatcgct	atgaagcggg	agatggcacg	1980
	tgaaggggga	ccgttgtgga	atccagtggt	accggtactt	ccggtactgg	ttatgtactc	2040
	aatttccaca	atgcagggga	ttccctgacg	atgacgatcc	aggctcccac	ggcaggcttg	2100
45	tacaatctta	caatcgata	ccgttctcct	catgatgaca	aacgcacgaa	tttctcttta	2160
	aatggcgaag	cgtctggaga	gctgggtact	tggaaaacgg	ctgattttta	agaaacgtcc	2220
	ggtggtaagg	ttctgttgaa	tgcaaggggc	aatcacgatc	gttttgaaac	aggctggggc	2280
	tggtatgata	tcgactacgt	caagctggag	ccagctgctg	accgtccacc	gcagtcggga	2340
	acgaaaacgc	tcataaatcc	gaatgcgaca	gtagaagcaa	aagcattgat	gaactacctg	2400
50	gttgatcaat	acgggaagaa	tatgctttcc	ggtcaagagg	atatgcccga	aattgatttg	2460
	cttcaagcga	atgtaggtaa	aaagccggct	attgcggcac	ttgacctgat	tgactattcc	2520
	ccaagcagag	cggaaacacg	tcttagttcc	acagagacgg	aaaaggcgat	tgaatgggat	2580
	aagcaagggg	gcattgttac	ctttgcatgg	cactggaaac	cgccgaaagg	tctgatcgat	2640
	acgcagggaa	aagaatggtg	gagaggcttc	tatgccgatt	cgactacatt	cgatatagaa	2700
55	tatgcgatga	atcatccaga	gtccgaagat	tataaattgc	ttattcgcca	catcgatgtg	2760
	attgcagggc	aattgaagaa	gttgaggat	gcgaaagtcc	ctgtcctgtt	ccgtcctttg	2820
	cacgaagcgg	agggcaaatg	gttctggtgg	ggcgctaaag	gtcctgagcc	tgttaaaaaa	2880
	ttgtatattt	tgatgcacga	tcgtttgact	aatgtgcaca	aattgaacaa	tctgatctgg	2940
	gtctggaaat	ctgttgctcc	cgactgggtat	ccgggagatg	agtatgtgga	tattttgagc	3000
60	ttcgactctt	atccgcgaagc	aggcgactac	agcccgcata	ttgcaaaaata	tgaagacctt	3060
	gttacatttg	gcaaggacaa	aaagctagtt	tgccatgagc	gaaaacggac	ctatcccggg	3120
	cccggtatctg	atgaaggcgt	atcaagccca	ttggagctgg	ttcgctacat	ggatggggga	3180
	ttctttgaga	cagcgcaaac	aaaacagctc	ttggagcatt	tgaaaaaaagt	gtataatcat	3240
	ccgaacgtca	ttacgcttga	aaagctcccg	actaacttaa	aaacgtatgg	cattaccgag	3300
65	caaccgtcag	taccgggcag	cttcacgctg	aacgcagcgg	gcgaaacggc	gaaagtaaag	3360
	ctgagctgga	cagcatcagc	gaatgcagca	agctatgaag	tgaagcggtc	gacgggtgaa	3420
	acggcgcggt	tcgccacagt	gcgagcggaa	gtatacggaa	cgagctacac	cgacacagcc	3480
	gtaacagcag	acacgacgta	ctattaccaa	gtcgtagcga	agaacgatgc	aggtcaaacg	3540
	gtttcgaaca	cggctagcgc	agcgcggaaa	gcggatactc	agcagccgac	aacgggattg	3600
70	gtgctccagt	atcgcacagc	ggatacaaat	gtgaacgaca	atcacttgaa	cccgcatctc	3660

```

caaatttttaa ataaaggtac aatctccgta cccgatcaacg agttgaaaat tccgtactac 3720
tacacgatcg acggtgaccg tgagcagaca ttcaactgcg actatgcggt gctgagctgc 3780
tcgaagctga atggttaagct ggttaaaatg gataaagctg caaccgggtgc tgattattat 3840
ttggaagtca gtttcaactc ggatgcaggc gtgttagcct ctggaggaag cacgggcgga 3900
5 attcaaaactc gtattcataa agcagactgg tcgaactata acgaaagtga cgattactcg 3960
tataaaggta cgcagacttc attcgacgat catacgaaag ctacgttgta tcacaatggc 4020
gtacttgttt ggggaaccga accgacagct aattaa 4056

```

```

10 <210> 4
    <211> 1350
    <212> PRT
    <213> Paenibacillus polymyxa

15 <400> 4
    Met Lys Ala Lys Asn Ser Ser Ser Ile Trp Ser Lys Arg Ser Lys Trp
      1          5          10          15

    Leu Pro Val Val Met Ala Cys Thr Ile Ile Val Gly Gly Ala Leu Pro
20      20          25          30

    Thr Pro Thr Val Val His Gly Gln Thr Ala Lys Thr Val Thr Ile Lys
      35          40          45

25 Val Asp Thr Ser Lys Asp Arg Lys Pro Ile Ser Pro Tyr Ile Tyr Gly
      50          55          60

    Thr Asn Gln Glu Leu Ala Gly Asp Glu Asn Leu Thr Ala Arg Arg Leu
      65          70          75          80

30 Gly Gly Asn Arg Met Thr Gly Tyr Asn Trp Glu Asn Asn Met Ser Asn
      85          90          95

    Ala Gly Ser Asp Trp Met Gln Ser Ser Asp Ser Tyr Leu Cys Asp Asn
35      100          105          110

    Ala Gly Leu Thr Lys Ala Glu Cys Glu Lys Pro Gly Ala Val Ala Thr
      115          120          125

40 Ser Phe His Asp Gln Ser Leu Lys Gln Gly Thr Tyr Ser Leu Val Thr
      130          135          140

    Leu Pro Met Ala Gly Tyr Val Ala Lys Asp Gly Asn Gly Ser Val Gln
      145          150          155          160

45 Glu Ser Glu Lys Ala Pro Ser Ala Arg Trp Asn Glu Val Val Asn Ala
      165          170          175

    Lys Asn Ala Pro Phe Gln Leu Gln Pro Asp Leu Lys Asp Asn Gln Val
50      180          185          190

    Tyr Ala Asp Glu Phe Val Asn Phe Leu Val Lys Lys Tyr Gly Val Ala
      195          200          205

55 Ser Thr Lys Thr Gly Val Lys Gly Tyr Ser Leu Asp Asn Glu Pro Ala
      210          215          220

    Leu Trp Ser His Thr His Pro Arg Ile His Gly Glu Lys Val Gly Ala
      225          230          235          240

60 Lys Glu Leu Val Asp Arg Ser Val Ser Leu Ser Lys Ala Ala Lys Ala
      245          250          255

    Val Asp Ala Gly Ala Glu Ile Phe Gly Pro Val Leu Tyr Gly Phe Gly
65      260          265          270

    Ala Tyr Lys Asp Leu Gln Thr Ala Pro Asp Trp Asn Ser Val Lys Gly
      275          280          285

70 Asn Tyr Ser Trp Phe Val Asp Tyr Tyr Leu Asp Gln Met Arg Leu Ser

```

	290	295	300
	Ser Gln Ala Glu Gly Lys Arg Leu Leu Asp Val Phe Asp Val His Trp		
	305	310	315 320
5	Tyr Pro Glu Ala Met Gly Gly Gly Ile Arg Ile Thr Asn Glu Val Gly		
	325	330	335
10	Asn Asp Glu Thr Lys Lys Ala Arg Met Gln Ala Pro Arg Thr Leu Trp		
	340	345	350
	Asp Pro Thr Tyr Lys Glu Asp Ser Trp Ile Ala Gln Trp Asn Ser Glu		
	355	360	365
15	Phe Leu Pro Leu Leu Pro Arg Leu Lys Gln Ser Val Asp Lys Tyr Tyr		
	370	375	380
	Pro Gly Thr Lys Leu Ala Leu Thr Glu Tyr Ser Tyr Gly Gly Glu Asn		
	385	390	395 400
20	Asp Ile Ser Gly Gly Ile Ala Met Ala Asp Val Leu Gly Ile Leu Gly		
	405	410	415
	Lys Asn Asp Val Tyr Met Ala Asn Tyr Trp Lys Leu Lys Asp Gly Ala		
25	420	425	430
	Asn Asn Tyr Val Ser Ala Ala Tyr Lys Leu Tyr Arg Asn Tyr Asp Gly		
	435	440	445
30	Lys Ser Ser Thr Phe Gly Asp Ile Ser Val His Ala Gln Thr Ser Asp		
	450	455	460
	Ile Val Asn Ser Ser Val His Ala Ser Val Thr Asp Ala Ser Tyr Lys		
	465	470	475 480
35	Glu Leu His Leu Val Val Met Asn Lys Ser Met Asp Ser Ala Phe Asp		
	485	490	495
	Ala Gln Phe Asp Leu Ser Gly Glu Thr Thr Tyr Gly Ser Gly Lys Val		
40	500	505	510
	Trp Gly Phe Asp Lys Asn Ser Ser Gln Ile Lys Glu Ala Ala Pro Ile		
	515	520	525
45	Thr Gln Ile Ser Gly Asn Arg Phe Thr Tyr Thr Val Pro Pro Leu Thr		
	530	535	540
	Ala Tyr His Ile Val Leu Thr Ala Gly Asn Asp Thr Pro Val Glu Asn		
	545	550	555 560
50	Pro Glu Ser Phe Ala Leu Arg Ala Glu Ala Gly Asp Gly Lys Ser Ile		
	565	570	575
	Tyr Leu Asp Ala Ser Ser Gly Val Val Gly Tyr Ser Val Gln Arg Ala		
55	580	585	590
	Thr Tyr Glu Asn Gly Pro Phe Ala Ala Val Ala Ser Asn Leu Val Glu		
	595	600	605
60	Thr Ser Tyr Thr Asp Thr Asn Val Thr Asn Gly Thr Ser Tyr Tyr Tyr		
	610	615	620
	Lys Ile Thr Ala Lys Thr Lys Thr Gly Thr Ser Ala Ser Asn Val Leu		
	625	630	635 640
65	Lys Ala Val Pro Arg Ala Pro Val Asp Gly Pro Asp Arg Tyr Glu Ala		
	645	650	655
	Glu Asp Gly Thr Leu Lys Gly Thr Val Val Glu Ser Ser Gly Thr Gly		
70	660	665	670

Phe Ser Gly Thr Gly Tyr Val Thr Asn Phe His Asn Ala Gly Asp Ser
 675 680 685
 5 Leu Thr Met Thr Ile Gln Ala Pro Thr Ala Gly Leu Tyr Asn Leu Thr
 690 695 700
 Ile Gly Tyr Arg Ser Pro His Asp Asp Lys Arg Thr Asn Phe Ser Leu
 705 710 715 720
 10 Asn Gly Lys Ala Ser Gly Glu Leu Val Leu Trp Lys Thr Ala Asp Phe
 725 730 735
 Lys Glu Thr Ser Gly Gly Lys Val Leu Leu Asn Ala Gly Ala Asn Thr
 15 740 745 750
 Ile Gly Phe Glu Thr Gly Trp Gly Trp Tyr Asp Ile Asp Tyr Val Lys
 755 760 765
 20 Leu Glu Pro Ala Ala Asp Arg Pro Pro His Ala Val Thr Lys Thr Leu
 770 775 780
 Ile Asn Pro Asn Ala Thr Val Glu Ala Lys Ala Leu Met Asn Tyr Leu
 785 790 795 800
 25 Val Asp Gln Tyr Gly Lys Asn Met Leu Ser Gly Gln Glu Asp Met Pro
 805 810 815
 Glu Ile Asp Trp Leu Gln Ala Asn Val Gly Lys Lys Pro Ala Ile Ala
 30 820 825 830
 Ala Leu Asp Leu Ile Asp Tyr Ser Pro Ser Arg Ala Glu His Gly Leu
 835 840 845
 35 Ser Ser Thr Glu Thr Glu Lys Ala Ile Glu Trp Asp Lys Gln Gly Gly
 850 855 860
 Ile Val Thr Phe Ala Trp His Trp Asn Ala Pro Lys Gly Leu Ile Asp
 865 870 875 880
 40 Thr Gln Gly Lys Glu Trp Trp Arg Gly Phe Tyr Ala Asp Ser Thr Thr
 885 890 895
 Phe Asp Ile Glu Tyr Ala Met Asn His Pro Glu Ser Glu Asp Tyr Lys
 45 900 905 910
 Leu Leu Ile Arg Asp Ile Asp Val Ile Ala Gly Gln Leu Lys Lys Leu
 915 920 925
 50 Gln Asp Ala Lys Val Pro Val Leu Phe Arg Pro Leu His Glu Ala Glu
 930 935 940
 Gly Lys Trp Phe Trp Trp Gly Ala Lys Gly Pro Glu Pro Val Lys Lys
 945 950 955 960
 55 Leu Tyr Ile Leu Met His Asp Arg Leu Thr Asn Val His Lys Leu Asn
 965 970 975
 Asn Leu Ile Trp Val Trp Asn Ser Val Ala Pro Asp Trp Tyr Pro Gly
 60 980 985 990
 Asp Glu Tyr Val Asp Ile Leu Ser Phe Asp Ser Tyr Pro Gln Ala Gly
 995 1000 1005
 65 Asp Tyr Ser Pro Gln Ile Ala Lys Tyr Glu Asp Leu Val Thr Leu Gly
 1010 1015 1020
 Lys Asp Lys Lys Leu Val Cys His Glu Arg Lys Arg Thr Tyr Pro Gly
 1025 1030 1035 1040
 70

Pro Gly Ser Asp Glu Gly Val Ser Ser Pro Leu Glu Leu Val Arg Tyr
 1045 1050 1055
 5 Met Val Trp Gly Phe Leu Glu Arg Arg Gln Thr Lys Gln Ser Leu Glu
 1060 1065 1070
 His Leu Lys Lys Val Tyr Asn His Pro Asn Val Ile Thr Leu Glu Lys
 1075 1080 1085
 10 Leu Pro Thr Asn Leu Lys Thr Tyr Gly Ile Thr Glu Gln Pro Ser Val
 1090 1095 1100
 Pro Gly Ser Phe Thr Leu Asn Ala Ala Gly Glu Thr Ala Lys Val Lys
 1105 1110 1115 1120
 15 Leu Ser Trp Thr Ala Ser Ala Asn Ala Ala Ser Tyr Glu Val Lys Arg
 1125 1130 1135
 Ser Thr Val Glu Asn Gly Ala Phe Ala Thr Val Ala Ser Asp Val Tyr
 1140 1145 1150
 20 Gly Ser Ser Tyr Thr Asp Thr Ala Val Thr Ala Asp Thr Thr Tyr Tyr
 1155 1160 1165
 Tyr Gln Val Val Ala Lys Asn Asp Ala Gly Gln Thr Val Ser Asn Thr
 1170 1175 1180
 Ala Ser Ala Ala Pro Lys Ala Asp Thr Gln Gln Pro Thr Thr Gly Leu
 1185 1190 1195 1200
 30 Val Leu Gln Tyr Arg Thr Ala Asp Thr Asn Val Asn Asp Asn His Leu
 1205 1210 1215
 Asn Pro His Phe Gln Ile Leu Asn Lys Gly Thr Ile Ser Val Pro Ile
 1220 1225 1230
 Asn Glu Leu Lys Ile Arg Tyr Tyr Tyr Thr Ile Asp Gly Asp Arg Glu
 1235 1240 1245
 40 Gln Thr Phe Asn Cys Asp Tyr Ala Val Leu Ser Cys Ser Lys Leu Asn
 1250 1255 1260
 Gly Lys Leu Val Lys Met Asp Lys Ala Ala Thr Gly Ala Asp Tyr Tyr
 1265 1270 1275 1280
 45 Leu Glu Val Ser Phe Asn Ser Asp Ala Gly Val Leu Ala Ser Gly Gly
 1285 1290 1295
 Ser Thr Gly Gly Ile Gln Thr Arg Ile His Lys Ala Asp Trp Ser Asn
 1300 1305 1310
 Tyr Asn Glu Ser Asp Asp Tyr Ser Tyr Lys Gly Thr Gln Thr Ser Phe
 1315 1320 1325
 55 Asp Asp His Thr Lys Ala Thr Leu Tyr His Asn Gly Val Leu Val Trp
 1330 1335 1340
 Gly Thr Glu Thr Ala Asn
 1345 1350
 60
 <210> 5
 <211> 2141
 <212> DNA
 65 <213> Paenibacillus pabuli
 <400> 5
 atgaaggcga aaaatagtag taatatatttg tccaaacggt ccaaattggct gcctgtcgctc 60
 atggcatgca cgattatagt agggggggct ctaccggctc caactgtagt tcacgggtcaa 120
 70 acggcaaaga ccgttaccat taaagtcgat acatccaagg atcgtaagcc tattagtcct 180

	tatatatacg	gtacgaattca	ggatttggcca	ggcgcgatgaaa	atctggctgc	cagacgactt	240
	ggtaggcaatc	gaatgaccgg	atacaactgg	gaaaaataata	tgtccaatgc	gggaagcgat	300
	tggcagcaat	ctagcgataa	cttttttatgc	aacaatgggtg	gcctgacaaa	agccgaatgt	360
5	gaaaggccgg	gagcagtgc	gacttcgttt	catgatcaat	cgctgaagct	gggcgcttat	420
	tctttagtca	cgctgccgac	ggccgggttat	gtggccaagg	atggaaaacgg	aagtgtgcag	480
	gaaagcgaac	aggctccttc	cgctcggttg	aatcagggtcg	taaatgccaa	aaatgcgccg	540
	ttccaactac	agcctgatct	gaatgacaat	caggtatatg	cggatgaatt	cgtaaathtt	600
	ttagtgaaaa	agtacggcgc	tgttcaaca	aaggcgggtg	tgaaggata	tgcgctcgac	660
	aatgaaccgc	ctctctggtc	gcatacgcat	ccgcgcattc	atggtgaaaa	agtcnnnnnn	720
10	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	780
	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnntcaaaact	gcacntgatt	840
	ggaacttctg	taaaaggcaa	ctatagctgg	ttcgtggact	attacctgga	tcaaagtgcg	900
	ctcaactcgc	aagccgaagg	caagagattg	ctggatgtat	tcgatgtgca	ctggatccc	960
	gaagcgatgg	gcggaggcat	acgaattaca	aatgaggtag	gcaatgacga	aacgaagaaa	1020
15	gccagaatgc	aggcgccctg	tacttttgtg	gacccgacct	acaagggaaga	tagctggatc	1080
	gctcaatgga	acagcgcat	cttgccctta	ctgcctcgat	tgaagcagtc	ggtggacaag	1140
	tattaccggg	gaaccaagct	ggcttttgacc	gagtatagct	acggcggcga	aaatgatatt	1200
	tccggcggtg	ttgctatgac	cgatgtgctg	ggcatcttgg	gcaaaaacga	cgtttatatg	1260
	gcgaactatt	ggaagttaa	ggatgggtgc	aacaactacg	ttagcgccgc	ttacaagctt	1320
20	taccgcaatt	atgacggaaa	aaacgctact	ttcggcgata	tcagcggtta	tgcgcaaacg	1380
	tccgatattg	ttaatgctc	ggtgcatgct	tccgtaacgg	atgcactcta	caaagaatcc	1440
	cacctcattg	tcatgaataa	aagcatggac	agcgcattcg	acgccccatt	cgatctttcc	1500
	ggcgagacga	cttacagttc	cggtaaaaaa	tggggcttcg	ataaaaaatag	ctcgcaaatt	1560
	aaggcaatga	gcaccaatcac	ggcaaatttca	ggcaaccgct	ttacctatac	agtaccacct	1620
25	ttgacggctt	atcacatcgt	gttgactgcc	gacaatgata	caactgtgcc	acctgtggaa	1680
	gatcctgaaa	gctttacgct	gagggctgag	gctggcgatg	ggaaagtoga	tttgtcctgg	1740
	gacgcttcca	gcggagttgt	gggttacagt	gtacagcggg	caacgtatga	aaacggctct	1800
	tttgctctg	tagcatccaa	cttgggtcgaa	acgtcttata	cggatacga	cgtaacgaac	1860
	ggcacttctt	actattataa	aataaccgca	aaaacaaagg	cgggaacgag	cgaatccaat	1920
30	gtcttgaaa	cggttccgcg	aacgcctgta	gacggcccgg	atcgctatga	agccgaagat	1980
	ggcactgcta	agggaaacct	tgtggaaatcc	agcgggacgc	gcttctccgg	tactggttat	2040
	gtaactaatt	tccacaatgc	aggggatccc	ctgacgatga	cgatccacta	gtgtcgacct	2100
	gcaggcgcg	gagctccagc	ttttgttccc	tttagtgagg	g	2141	
35	<210>	6					
	<211>	695					
	<212>	PRT					
	<213>	Paenibacillus pabuli					
40	<400>	6					
	Met Lys Ala	Lys Asn Ser	Ser Asn Ile	Leu Ser Lys	Arg Ser Lys	Trp	
	1	5	10	'	15		
45	Leu Pro Val	Val Met Ala	Cys Thr Ile	Ile Val Gly	Gly Ala Leu	Pro	
	20	25	30				
	Ala Pro Thr	Val Val His	Gly Gln Thr	Ala Lys Thr	Val Thr Ile	Lys	
	35	40	45				
50	Val Asp Thr	Ser Lys Asp	Arg Lys Pro	Ile Ser Pro	Tyr Ile Tyr	Gly	
	50	55	60				
	Thr Asn Gln	Asp Leu Ala	Gly Asp Glu	Asn Leu Ala	Ala Arg Arg	Leu	
55	65	70	75	80			
	Gly Gly Asn	Arg Met Thr	Gly Tyr Asn	Trp Glu Asn	Asn Met Ser	Asn	
	85	90	95				
60	Ala Gly Ser	Asp Trp Gln	Gln Ser Ser	Asp Asn Phe	Leu Cys Asn	Asn	
	100	105	110				
	Gly Gly Leu	Thr Lys Ala	Glu Cys Glu	Lys Pro Gly	Ala Val Thr	Thr	
	115	120	125				
65	Ser Phe His	Asp Gln Ser	Leu Lys Leu	Gly Ala Tyr	Ser Leu Val	Thr	
	130	135	140				
	Leu Pro Met	Ala Gly Tyr	Val Ala Lys	Asp Gly Asn	Gly Ser Val	Gln	
70	145	150	155	160			

Glu Ser Glu Gln Ala Pro Ser Ala Arg Trp Asn Gln Val Val Asn Ala
 165 170 175
 5 Lys Asn Ala Pro Phe Gln Leu Gln Pro Asp Leu Asn Asp Asn Gln Val
 180 185 190
 Tyr Ala Asp Glu Phe Val Asn Phe Leu Val Lys Lys Tyr Gly Ala Ala
 195 200 205
 10 Ser Thr Lys Ala Gly Val Lys Gly Tyr Ala Leu Asp Asn Glu Pro Ala
 210 215 220
 Leu Trp Ser His Thr His Pro Arg Ile His Gly Glu Lys Val Xaa Xaa
 225 230 235 240
 Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
 245 250 255
 20 Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
 260 265 270
 Xaa Xaa Phe Lys Leu His Xaa Ile Gly Thr Ser Val Lys Gly Asn Tyr
 275 280 285
 25 Ser Trp Phe Val Asp Tyr Tyr Leu Asp Gln Met Arg Leu Asn Ser Gln
 290 295 300
 Ala Glu Gly Lys Arg Leu Leu Asp Val Phe Asp Val His Trp Tyr Pro
 305 310 315 320
 Glu Ala Met Gly Gly Gly Ile Arg Ile Thr Asn Glu Val Gly Asn Asp
 325 330 335
 35 Glu Thr Lys Lys Ala Arg Met Gln Ala Pro Arg Thr Leu Trp Asp Pro
 340 345 350
 Thr Tyr Lys Glu Asp Ser Trp Ile Ala Gln Trp Asn Ser Ala Phe Leu
 355 360 365
 40 Pro Leu Leu Pro Arg Leu Lys Gln Ser Val Asp Lys Tyr Tyr Pro Gly
 370 375 380
 Thr Lys Leu Ala Leu Thr Glu Tyr Ser Tyr Gly Gly Glu Asn Asp Ile
 385 390 395 400
 Ser Gly Gly Ile Ala Met Thr Asp Val Leu Gly Ile Leu Gly Lys Asn
 405 410 415
 50 Asp Val Tyr Met Ala Asn Tyr Trp Lys Leu Lys Asp Gly Ala Asn Asn
 420 425 430
 Tyr Val Ser Ala Ala Tyr Lys Leu Tyr Arg Asn Tyr Asp Gly Lys Asn
 435 440 445
 55 Ala Thr Phe Gly Asp Ile Ser Val Asn Ala Gln Thr Ser Asp Ile Val
 450 455 460
 Asn Ser Ser Val His Ala Ser Val Thr Asp Ala Ser Tyr Lys Glu Leu
 465 470 475 480
 His Leu Ile Val Met Asn Lys Ser Met Asp Ser Ala Phe Asp Ala Gln
 485 490 495
 65 Phe Asp Leu Ser Gly Glu Thr Thr Tyr Ser Ser Gly Lys Ile Trp Gly
 500 505 510
 Phe Asp Lys Asn Ser Ser Gln Ile Lys Ala Val Ala Pro Ile Thr Gln
 515 520 525
 70

Ile Ser Gly Asn Arg Phe Thr Tyr Thr Val Pro Pro Leu Thr Ala Tyr
 530 535 540
 His Ile Val Leu Thr Ala Asp Asn Asp Thr Pro Val Pro Pro Val Glu
 5 545 550 555 560
 Asp Pro Glu Ser Phe Thr Leu Arg Ala Glu Ala Gly Asp Gly Lys Val
 565 570 575
 10 Asp Leu Ser Trp Asp Ala Ser Ser Gly Val Val Gly Tyr Ser Val Gln
 580 585 590
 Arg Ala Thr Tyr Glu Asn Gly Pro Phe Ala Ala Val Ala Ser Asn Leu
 595 600 605
 15 Val Glu Thr Ser Tyr Thr Asp Thr Asn Val Thr Asn Gly Thr Ser Tyr
 610 615 620
 Tyr Tyr Lys Ile Thr Ala Lys Thr Lys Ala Gly Thr Ser Glu Ser Asn
 20 625 630 635 640
 Val Leu Lys Ala Val Pro Arg Thr Pro Val Asp Gly Pro Asp Arg Tyr
 645 650 655
 25 Glu Ala Glu Asp Gly Thr Leu Lys Gly Thr Ile Val Glu Ser Ser Gly
 660 665 670
 Thr Gly Phe Ser Gly Thr Gly Tyr Val Thr Asn Phe His Asn Ala Gly
 675 680 685
 30 Asp Ser Leu Thr Met Thr Ile
 690 695
 35

CLAIMS

1. A xyloglucanase enzyme belonging to family 44 of
5 glycosylhydrolases, which enzyme exhibits a relative
xyloglucanase activity of at least 30% between pH 5 and pH 8.
2. The enzyme according to claim 1, which is obtainable from a
microorganism, preferably from a bacterium or a fungus,
10 including a yeast.
3. The enzyme according to claim 2, which is obtained or
obtainable from a bacterium, preferably from a Gram positive
bacterium, preferably from the *Bacillus/Lactobacillus*
15 subdivision, preferably a species from the genus *Paenibacillus*,
especially *Paenibacillus polymyxa*, e.g. *Paenibacillus polymyxa*,
ATCC 832.
4. The enzyme according to claim 2, which is derived or
20 derivable from a bacterium, preferably from a Gram positive
bacterium, preferably from the *Bacillus/Lactobacillus*
subdivision, preferably a species from the genus *Paenibacillus*,
preferably from the strain *Paenibacillus sp.*, DSM 13329.
- 25 5. A family 44 xyloglucanase which is selected from one of
(a) a polypeptide encoded by the DNA sequence of positions 121-
1677 of SEQ ID NO:1;
(b) a polypeptide produced by culturing a cell comprising the
sequence of SEQ ID NO:1 under conditions wherein the DNA
30 sequence is expressed;
(c) a xyloglucanase enzyme having a sequence of at least 60%
identity to positions 40-559 of SEQ ID NO:2 when identity is
determined by GAP provided in the GCG program package using a
GAP creation penalty of 3.0 and GAP extension penalty of 0.1; or
35 (d) a polypeptide encoded by the xyloglucanase encoding part of
the DNA sequence obtainable from the plasmid in *Escherichia coli*
DSM 13321.
6. The xyloglucanase according to claim 5 which is obtained or

obtainable from a bacterium, preferably from a Gram positive bacterium, preferably from the *Bacillus/Lactobacillus* subdivision, preferably a species from the genus *Paenibacillus*, especially a *Paenibacillus polymyxa*, e.g. *Paenibacillus*

5 *polymyxa*, ?.

7. The family 44 xyloglucanase according to claim 5, which is selected from one of

(a) a polypeptide encoded by the DNA sequence of positions 121-
10 1677 of SEQ ID NO:3;

(b) a polypeptide produced by culturing a cell comprising the sequence of SEQ ID NO:3 under conditions wherein the DNA sequence is expressed; or

(c) a polypeptide encoded by the xyloglucanase encoding part of
15 the DNA sequence obtainable from the plasmid in *Escherichia coli* DSM 13322.

8. The xyloglucanase according to claim 7 which is obtained or obtainable from a bacterium, preferably from a Gram positive
20 bacterium, preferably from the *Bacillus/Lactobacillus* subdivision, preferably a species from the genus *Paenibacillus*, especially a *Paenibacillus polymyxa*, e.g. *Paenibacillus polymyxa*, ATCC 832.

25 9. The family 44 xyloglucanase according to claim 5, which is selected from one of

(a) a polypeptide encoded by the DNA sequence of positions 121-
1677 of SEQ ID NO:5;

(b) a polypeptide produced by culturing a cell comprising the
30 sequence of SEQ ID NO:5 under conditions wherein the DNA sequence is expressed; or

(c) a polypeptide encoded by the xyloglucanase encoding part of the DNA sequence obtainable from the plasmid in *Escherichia coli* DSM 13323.

35

10. The xyloglucanase according to claim 9 which is obtained or obtainable from a bacterium, preferably a strain from the genus *Paenibacillus*, especially a strain of *Paenibacillus sp.*, DSM 13329.

11. An isolated xyloglucanase enzyme, in which the enzyme is (i) free from homologous impurities, and (ii) produced by culturing a cell comprising the DNA sequence of positions 121-1677 of SEQ ID NO:1, 3 or 5, wherein the enzyme is produced and isolated.
12. An enzyme preparation comprising the enzyme according to any of the claims 1, 5 or 11.
13. The preparation according to claim 12 which further comprises one or more enzymes selected from the group consisting of proteases, cellulases (endoglucanases), β -glucanases, hemicellulases, lipases, peroxidases, laccases, α -amylases, glucoamylases, cutinases, pectinases, reductases, oxidases, phenoloxidases, ligninases, pullulanases, pectate lyases, xyloglucanases, xylanases, pectin acetyl esterases, polygalacturonases, rhamnogalacturonases, pectin lyases, other mannanases, pectin methylesterases, cellobiohydrolases, transglutaminases; or mixtures thereof.
14. An isolated polynucleotide molecule encoding a polypeptide having xyloglucanase activity which polynucleotide molecule hybridizes to a denatured double-stranded DNA probe under medium stringency conditions, wherein the probe is selected from the group consisting of DNA probes comprising the sequence shown in positions 121-1677 of SEQ ID NO:1, 3 or 5, and DNA probes comprising a subsequence of positions 121-1677 of SEQ ID NO:1, 3 or 5, the subsequence having a length of at least about 100 base pairs.
16. An expression vector comprising the following operably linked elements: a transcription promoter; a DNA segment selected from the group consisting of (a) polynucleotide molecules encoding a polypeptide having xyloglucanase activity comprising a nucleotide sequence as shown in SEQ ID NO:1, 3 or 5 from nucleotide 121 to nucleotide 1677, (b) polynucleotide molecules encoding a polypeptide having xyloglucanase activity that is at least 60% identical to the amino acid sequence of SEQ

ID NO:2 from amino acid residue 40 to amino acid residue 559, and (c) degenerate nucleotide sequences of (a) or (b); and a transcription terminator.

- 5 17. A cultured cell into which has been introduced an expression vector according to claim 16, wherein said cell expresses the polypeptide encoded by the DNA segment.
- 10 18. A method of producing a polypeptide having xyloglucanase activity comprising culturing a cell into which has been introduced an expression vector according to claim 16, whereby said cell expresses a polypeptide encoded by the DNA segment; and recovering the polypeptide.
- 15 19. A detergent composition comprising the enzyme preparation according to claim 12 or the enzyme according to any of the claims 1, 5 and 11.
- 20 20. A process for machine treatment of fabrics which process comprises treating fabric during a washing cycle of a machine washing process with a washing solution containing the enzyme preparation according to claim 12 or the enzyme according to any of the claims 1, 5 and 11.
- 25 21. Use of the enzyme preparation according to claim 12 or the enzyme according to any of the claims 1, 5 and 11 in the textile industry for improving the properties of cellulosic fibres, yarn, woven or non-woven fabric.
- 30 22. The use according to claim 21, wherein the enzyme preparation or the enzyme is used in a scouring process step.
- 35 23. Use of the enzyme preparation according to claim 12 or the enzyme according to any of the claims 1, 5 and 11 in the cellulose fiber processing industry for ratting of fibers selected from the group consisting of hemp, jute, flax and linen.

ABSTRACT

FAMILY 44 XYLOGLUCANASES

5
Xyloglucanases belonging to family 44 of glycosylhydrolases and
having a relative xyloglucanase activity of at least 30% between
pH 5 and pH 8 are derived from the genus *Paenibacillus*,
especially a strain of *Paenibacillus polymyxa* or *Paenibacillus*
10 *sp.* The xyloglucanases show high performance in conventional
detergent compositions.